


PB82-132994

Pesticide Registration Standard: Carboxin

(U.S.) Environmental Protection Agency
Washington, DC

Aug 81



U.S. Department of Commerce
National Technical Information Service

NTIS

REPORT DOCUMENTATION PAGE	1. REPORT NO. EPA 540/RS-81-007	2.	3. Recipient's Accession No. PB82 132994
4. Title and Subtitle Pesticide Registration Standard Carboxin			5. Report Date August, 1981
7. Author(s) USEPA, Special Pesticide Review Division (SPRD) and Registration Division (RD)			6. 335021
9. Performing Organization Name and Address U.S. Environmental Protection Agency 401 M Street, S.W. Washington, D.C. 20460			8. Performing Organization Rept. No.
12. Sponsoring Organization Name and Address Same as # 9			10. Project/Task/Work Unit No.
15. Supplementary Notes			11. Contract(C) or Grant(G) No. (C) (G)
16. Abstract (Limit: 200 words) This document contains information regarding reregistration of pesticide products containing Carboxin, including how to register under the registration standard, regulatory position and rationale, and summaries of data requirements and data gaps. Also include is a bibliography containing citations of all studies reviewed by EPA in arriving at the positions and conclusions contained in the standard.			13. Type of Report & Period Covered
17. Document Analysis a. Descriptors Registration Standard Carboxin b. Identifiers/Open-Ended Terms c. COSATI Field/Group			14.
18. Availability Statement:		19. Security Class (This Report) Unclassified	21. No. of Pages 120
		20. Security Class (This Page)	22. Price

United States
Environmental Protection
Agency

Office of
Pesticides and Toxic Substances
Washington DC 20460

August 1981


P882-132994

Pesticides



5,6-dihydro-2-methyl-1,4- oxathiin-3-carboxanilide (Carboxin)

Pesticide Registration Standard



REPRODUCED BY
NATIONAL TECHNICAL
INFORMATION SERVICE
U. S. DEPARTMENT OF COMMERCE
SPRINGFIELD, VA. 22161

CARBOXIN

Pesticide Registration Standard

Bruce Kapner	Project Manager (SPRD)
Alex Arce	Toxicologist (HED)
Jolene Chinchilli	Project Manager (SPRD)
Ed Fite	Wildlife Biologist (HED)
Linda Garczynski	Writer/Editor (SPRD)
Robert Holst	Plant Physiologist (HED)
Steve Hopkins	Plant Physiologist (HED)
Sam Howard	Environmental Chemist (HED)
Ray Kent	Chemist (HED)
Greg Weidemann	Plant Pathologist (BFSD)
Gene Wilson	Product Manager (RD)
Linda Zygadlo	Economist (BFSD)

August 1981

Office of Pesticides and Toxic Substances

Environmental Protection Agency
401 M Street, S.W.
Washington, D.C. 20460

- TABLE OF CONTENTS -

	Page
Chapter I How to Register under a Registration Standard.....	1
Chapter II Regulatory Position and Rationale.....	9
Chapter III Summary of Data Requirements and Data Gaps.....	17
Chapter IV Product Chemistry.....	29
Chapter V Environmental Fate.....	33
Chapter VI Toxicology.....	41
Chapter VII Residue Chemistry.....	47
Chapter VIII Ecological Effects.....	55
Chapter IX Bibliography.....	58

I. HOW TO REGISTER UNDER A REGISTRATION STANDARD

- A. Organization of the Standard
- B. Purpose of the Standard
- C. Requirement to Reregister Under the Standard
- D. "Product Specific" Data and "Generic" Data
- E. Data Compensation Requirements under FIFRA 3(c)(1)(D)
- F. Obtaining Data to Fill "Data Gaps"; FIFRA 3(c)(2)(B)
- F. Amendments to the Standard

A. ORGANIZATION OF THE STANDARD

This first chapter explains the purpose of a Registration Standard and summarizes the legal principles involved in registering or reregistering under a Standard. The second chapter sets forth the requirements that must be met to obtain or retain registration for products covered by this particular Registration Standard. In the remaining chapters, the Agency reviews the available data by scientific discipline, discusses the Agency's concerns with the identified potential hazards, and logically develops the conditions and requirements that would reduce those hazards to acceptable levels.

B. PURPOSE OF THE STANDARD

Section 3 of the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) provides that "no person in any State may distribute, sell, offer for sale, hold for sale, ship, deliver for shipment, or receive (and having so received) deliver or offer to deliver, to any person any pesticide which is not registered with the Administrator [of EPA]." To approve the registration of a pesticide, the Administrator must find, pursuant to Section 3(c)(5) that:

- "(1) its composition is such as to warrant the proposed claims for it;
- (2) its labeling and other material required to be submitted comply with the requirements of this Act;
- (3) it will perform its intended function without unreasonable adverse effects on the environment; and
- (4) when used in accordance with widespread and commonly recognized practice it will not generally cause unreasonable adverse effects on the environment."

In making these findings, the Agency reviews a wide range of data which registrants are required to submit, and assesses the risks and benefits associated with the use of the proposed pesticide. However, the established approach to making these findings has been found to be defective on two counts.

First, EPA and its predecessor agency, the United States Department of Agriculture (USDA), routinely reviewed registration applications on a "product by product" basis, evaluating each product-specific application somewhat independently. In the review of products containing similar components, there was little opportunity for a retrospective review of the full range of

pertinent data available in Agency files and in the public literature. Thus the "product by product" approach was often inefficient and sometimes resulted in inconsistent or incomplete regulatory judgments.

Second, over the years, as a result of inevitable and continuing advances in scientific knowledge, methodology, and policy, the data base for many pesticides came to be considered inadequate by current scientific and regulatory standards. Given the long history of pesticide regulation in several agencies, it is even likely that materials may have been lost from the data files. When EPA issued new requirements for registration in 1975 (40 CFR 162) and proposed new guidelines for hazard testing in 1978 (43 FR 29686, July 10, 1978 and 43 FR 37336, August 22, 1978), many products that had already been registered for years were being sold and used without the same assurances of human and environmental safety as was being required for new products. Because of this inconsistency, Congress directed EPA to reregister all previously registered products, so as to bring their registrations and their data bases into compliance with current requirements [See FIFRA Section 3(g)].

Facing the enormous job of re-reviewing and calling-in new data for the approximately 35,000 current registrations, and realizing the inefficiencies of the "product by product" approach, the Agency decided that a new, more effective method of review was needed.

A new review procedure has been developed. Under it, EPA publishes documents called Registration Standards, each of which discusses a particular pesticide active ingredient. Each Registration Standard summarizes all the data available to the Agency on a particular active ingredient and its current uses, and sets forth the Agency's comprehensive position on the conditions and requirements for registration of all existing and future products which contain that active ingredient. These conditions and requirements, all of which must be met to obtain or retain full registration or reregistration under Section 3(c)(5) of FIFRA, include the submission of needed scientific data which the Agency does not now have, compliance with standards of toxicity, composition, labeling, and packaging, and satisfaction of the data compensation provisions of FIFRA Section 3(c)(1)(D).

The Standard will also serve as a tool for product classification. As part of the registration of a pesticide product, EPA may classify each product for "general use" or "restricted use" [FIFRA Section 3(d)]. A pesticide is classified for "restricted use" when some special regulatory restriction is needed to ensure against unreasonable adverse effects to man or the environment. Many such risks of unreasonable adverse effects can be lessened if expressly-designed label precautions are strictly followed. Thus the special regulatory restriction for a "restricted use" pesticide is usually a requirement that it be applied only by, or under the supervision of, an applicator who has been certified by the State or Federal government as being competent to use the pesticide safely, responsibly, and in accordance with label directions. A restricted-use pesticide can have other regulatory restrictions [40 CFR 162.11(c)(5)] instead of, or in addition to, the certified applicator requirement. These other regulatory restrictions may include such actions as seasonal or regional limitations on use, or a requirement for the monitoring of residue levels after use. A pesticide classified for "general

use," or not classified at all, is available for use by any individual who is in compliance with State or local regulations. The Registration Standard review compares information about potential adverse effects of specific uses of the pesticide with risk criteria listed in 40 CFR 162.11(c), and thereby determines whether a product needs to be classified for "restricted use." If the Standard does classify a pesticide for "restricted use," this determination is stated in the second chapter.

C. REQUIREMENT TO REREGISTER UNDER THE STANDARD

FIFRA Section 3(g), as amended in 1978, directs EPA to reregister all currently registered products as expeditiously as possible. Congress also agreed that reregistration should be accomplished by the use of Registration Standards.

Each registrant of a currently registered product to which this Standard applies, and who wishes to continue to sell or distribute his product in commerce, must apply for reregistration. His application must contain proposed labeling that complies with this Standard.

EPA will issue a notice of intent to cancel the registration of any currently registered product to which this Standard applies if the registrant fails to comply with the procedures for reregistration set forth in the Guidance Package which accompanies this Standard.

D. "PRODUCT SPECIFIC" DATA AND "GENERIC" DATA

In the course of developing this Standard, EPA has determined the types of data needed for evaluation of the properties and effects of products to which the Standard applies, in the disciplinary areas of Product Chemistry, Environmental Fate, Toxicology, Residue Chemistry, and Ecological Effects. These determinations are based primarily on the data Guidelines proposed in 43 FR 29696, July 10, 1978; 43 FR 37336, August 22, 1978; and 45 FR 72948, November 3, 1980, as applied to the use patterns of the products to which this Standard applies. Where it appeared that data from a normally applicable Guidelines requirement was actually unnecessary to evaluate these products, the Standard indicates that the requirement has been waived. On the other hand, in some cases studies not required by the Guidelines may be needed because of the particular composition or use pattern of products the Standard covers; if so, the Standard explains the Agency's reasoning. Data guidelines have not yet been proposed for the Residue Chemistry discipline, but the requirements for such data have been in effect for some time and are, the Agency believes, relatively familiar to registrants. Data which we have found are needed to evaluate the registrability of some products covered by the Standard may not be needed for the evaluation of other products, depending upon the composition, formulation type, and intended uses of the product in question. The Standard states which data requirements apply to which product categories. (See the third chapter.) The various kinds of data normally required for registration of a pesticide product can be divided into two basic groups:

1. Data that are product specific, i.e. data that relate only to the properties or effects of a product with a particular composition (or a group of products with closely similar composition); and
2. Generic data that pertains to the properties or effects of a particular ingredient, and thus are relevant to an evaluation of the risks and benefits of all products containing that ingredient (or all such products having a certain use pattern), regardless of any such product's unique composition.

The Agency requires certain "product specific" data for each product to characterize the product's particular composition and physical/chemical properties (Product Chemistry), and to characterize the product's acute toxicity (which is a function of its total composition). The applicant for registration or reregistration of any product, whether it is a manufacturing-use or end-use product, and without regard to its intended use pattern, must submit or cite enough of this kind of data to allow EPA to evaluate the product. For such purposes, "product specific" data on any product other than the applicant's is irrelevant, unless the other product is closely similar in composition to the applicant's. (Where it has been found practicable to group similar products for purposes of evaluating, with a single set of tests, all products in the group, the Standard so indicates.) "Product specific" data on the efficacy of particular end-use products are also required where the exact formulation may affect efficacy and where failure of efficacy could cause public health problems.

All other data needed to evaluate pesticide products concern the properties or effects of a particular ingredient of products (normally a pesticidally active ingredient, but in some cases a pesticidally inactive, or "inert", ingredient). Some data in this "generic" category are required to evaluate the properties and effects of all products containing that ingredient [e.g., the acute LD-50 of the active ingredient in its technical or purer grade; see proposed guidelines, 43 FR 37355].

Other "generic" data are required to evaluate all products which both contain a particular ingredient and are intended for certain uses (see, e.g., proposed guidelines, 43 FR 37363, which requires subchronic oral testing of the active ingredient with respect to certain use patterns only). Where a particular data requirement is use-pattern dependent, it will apply to each end-use product which is to be labeled for that use pattern (except where such end-use product is formulated from a registered manufacturing-use product permitting such formulations) and to each manufacturing-use product with labeling that allows it to be used to make end-use products with that use pattern. Thus, for example, a subchronic oral dosing study is needed to evaluate the safety of any manufacturing-use product that legally could be used to make an end-use, food-crop pesticide. But if an end-use product's label specified it was for use only in ways that involved no food/feed exposure and no repeated human exposure, the subchronic oral dosing study would not be

required to evaluate the product's safety; and if a manufacturing-use product's label states that the product is for use only in making end-use products not involving food/feed use or repeated human exposure, that subchronic oral study would not be relevant to the evaluation of the manufacturing-use product either.

If a registrant of a currently registered manufacturing-use or end-use product wishes to avoid the costs of data compensation [under FIFRA Section 3(c)(1)(D)] or data generation [under Section 3(c)(2)(B)] for "generic" data that is required only with respect to some use patterns, he may elect to delete those use patterns from his labeling at the time he reregisters his product. An applicant for registration of a new product under this Standard may similarly request approval for only certain use patterns.

E. DATA COMPENSATION REQUIREMENTS UNDER FIFRA 3(c)(1)(D)

Under FIFRA Section 3(c)(1)(D), an applicant for registration, reregistration, or amended registration must offer to pay compensation for certain existing data the Agency has used in developing the Registration Standard. The data for which compensation must be offered are all data which are described by all of the following criteria:

1. The data were first submitted to EPA (or to its predecessor agencies, USDA or FDA), on or after January 1, 1970;
2. The data were submitted to EPA (or USDA or FDA) by some other applicant or registrant in support of an application for an experimental use permit, an amendment adding a new use to a registration, or for registration, or to support or maintain an existing registration;
3. They are the kind of data which are relevant to the Agency's decision to register or reregister the applicant's product under the Registration Standard, taking into account the applicant's product's composition and intended use pattern(s);
4. The Agency has found the data to be valid and usable in reaching regulatory conclusions; and
5. They are not data for which the applicant has been exempted by FIFRA Section 3(c)(2)(D) from the duty to offer to pay compensation. (This exemption applies to the "generic" data concerning the safety of an active ingredient of the applicant's product, not to "product specific" data. The exemption is available only to applicants whose product is labeled for end-uses for which the active ingredient in question is present in the applicant's product because of his use of another registered product containing that active ingredient which he purchases from another producer.

An applicant for reregistration of an already registered product under this Standard, or for registration of a new product under this Standard, accordingly must determine which of the data used by EPA in developing the Standard must be

the subject of an offer to pay compensation, and must submit with his application the appropriate statements evidencing his compliance with FIFRA Section 3(c)(1)(D).

An applicant would never be required to offer to pay for "product specific" data submitted by another firm. In many, if not in most cases, data which are specific to another firm's product will not suffice to allow EPA to evaluate the applicant's product, that is, will not be useful to the Agency in determining whether the applicant's product is registrable. There may be cases, however, where because of close similarities between the composition of two or more products, another firm's data may suffice to allow EPA to evaluate some or all of the "product specific" aspects of the applicant's product. In such a case, the applicant may choose to cite that data instead of submitting data from tests on his own product, and if he chooses that option, he would have to comply with the offer-to-pay requirements of Section 3(C)(1)(D) for that data.

Each applicant for registration or reregistration of a manufacturing-use product, and each applicant for registration or reregistration of an end-use product, who is not exempted by FIFRA Section 3(c)(2)(D), must comply with the Section 3(c)(1)(D) requirements with respect to each item of "generic" data that relates to his product's intended uses.

A detailed description of the procedures an applicant must follow in applying for reregistration (or new registration) under this Standard is found in the Guidance Package for this Standard.

F. OBTAINING DATA TO FILL "DATA GAPS"; FIFRA 3(c)(2)(B)

Some of the kinds of data EPA needs for its evaluation of the properties and effects of products to which this Standard applies have never been submitted to the Agency (or, if submitted, have been found to have deficiencies rendering them inadequate for making registrability decisions) and have not been located in the published literature search that EPA conducted as part of preparing this Standard. Such instances of missing but required data are referred to in the Standard as "data gaps".

FIFRA Section 3(c)(2)(B), added to FIFRA by the Congress in 1978, authorizes EPA to require registrants to whom a data requirement applies to generate (or otherwise produce) data to fill such "gaps" and submit those data to EPA. EPA must allow a reasonably sufficient period for this to be accomplished. If a registrant fails to take appropriate and timely steps to fill the data gaps identified by a section 3(c)(2)(B) order, his product's registration may be suspended until the data are submitted. A mechanism is provided whereby two or more registrants may agree to share in the costs of producing data for which they are both responsible.

The Standard lists, in the third chapter, the "generic" data gaps and notes the classes of products to which these data gaps pertain. The Standard also points out that to be registrable under the Standard, a product must be supported by certain required "product specific" data. In some cases, the Agency may possess sufficient "product specific" data on one currently registered product,

but may lack such data on another. Only those Standards which apply to a very small number of currently registered products will attempt to state definitively the "product specific" data gaps on a "product by product" basis. (Although the Standard will in some cases note which data that EPA does possess would suffice to satisfy certain "product specific" data requirements for a category of products with closely similar composition characteristics.)

As part of the process of reregistering currently registered products, EPA will issue Section 3(c)(2)(B) directives requiring the registrants to take appropriate steps to fill all identified data gaps -- whether the data in question are "product specific" or "generic" -- in accordance with a schedule.

Persons who wish to obtain registrations for new products under this Standard will be required to submit (or cite) sufficient "product specific" data before their applications are approved. Upon registration, they will be required under Section 3(c)(2)(B) to take appropriate steps to submit data needed to fill "generic" data gaps. (We expect they will respond to this requirement by entering into cost-sharing agreements with other registrants who previously have been told they must furnish the data.) The Guidance Package for this Standard details the steps that must be taken by registrants to comply with Section 3(c)(2)(B).

G. AMENDMENTS TO THE STANDARD

Applications for registration which propose uses or formulations that are not presently covered by the Standard, or which present product compositions, product chemistry data, hazard data, toxicity levels, or labeling that do not meet the requirements of the Standard, will automatically be considered by the Agency to be requests for amendments to the Standard. In response to such applications, the Agency may request additional data to support the proposed amendment to the Standard, or may deny the application for registration on the grounds that the proposed product would cause unreasonable adverse effects to the environment. In the former case, when additional data have been satisfactorily supplied, and providing that the data do not indicate the potential for unreasonable adverse effects, the Agency will then amend the Standard to cover the new registration.

Each Registration Standard is based upon all data and information available to the Agency's reviewers on a particular date prior to the publication date. This "cut-off" date is stated at the beginning of the second chapter. Any subsequent data submissions and any approved amendments will be incorporated into the Registration Standard by means of addenda, which are available for inspection at EPA in Washington, D.C., or copies of which may be requested from the Agency. When all the present "data gaps" have been filled and the submitted data have been reviewed, the Agency will revise the Registration Standard. Thereafter, when the Agency determines that the internally maintained addenda have significantly altered the conditions for registration under the Standard, the document will be updated and re-issued.

While the Registration Standard discusses only the uses and hazards of products containing the designated active ingredient(s), the Agency is also concerned with the potential hazards of some inert ingredients and impurities.

Independent of the development of any one Standard, the Agency has initiated the evaluation of some inert pesticide ingredients. Where the Agency has identified inert ingredients of concern in a specific product to which the Standard applies, these ingredients will be pointed out in the Guidance Package.

II. REGULATORY POSITION AND RATIONALE

- A. Introduction
- B. Description of Chemical
- C. Regulatory Position
- D. Regulatory Rationale
- E. Criteria for Registration Under the Standard
- F. Acceptable Ranges and Limits
- G. Required Labeling
- H. Tolerance Reassessment


A. INTRODUCTION

This chapter presents the Agency's regulatory position and rationale based on an evaluation of all registered products containing carboxin as the sole active ingredient. After briefly describing carboxin, this chapter presents the regulatory position and rationale, and the criteria for registration of products containing this chemical. These include labeling considerations, and tolerance reassessment. A summary of data requirements is contained in Chapter III. Data supporting this regulatory position are discussed in each of the disciplinary chapters, IV through VIII. Also considered in this Standard are Special Local Need (SLN) registrations which are issued by a State in accordance with FIFRA section 24(c). Valid State registrations will be subject to reregistration under the EPA Registration Standard System, in the same manner as registration issued under section 3 of FIFRA.

B. DESCRIPTION OF CHEMICAL

Carboxin [5,6-dihydro-2-methyl-1,4-oxathiin-3-carboxanilide] is a systemic fungicide registered for use as a seed treatment to protect barley, corn, wheat, oats, cotton, and peanuts; and registered for SLN use only in the states of Alabama, Georgia, North Carolina, Oklahoma, South Carolina, and Texas as a combined foliar and soil application on peanut plants to control Southern blight.

The structural formula is:



Uniroyal, Inc. is the principal producer and registrant. Carboxin (technical material) is formulated into a formulation intermediate, a wettable powder/dust, a dust, two ready-to-use liquids (flowable), two soluble concentrate liquids; and also a granular formulation and flowable concentrate for SLN use only.

Carboxin is unusual because it is systemically toxic to phytopathogenic Basidiomycetes. It is less toxic to phytopathogenic fungi of other classes, therefore, it is commonly applied in combination with thiram or

captan, or applied to seed previously treated with other fungicides to broaden the spectrum of diseases controlled. Any combination (i.e. mixture) of carboxin with other pesticides will not be considered in this Standard at this time.

C. REGULATORY POSITION

Based on a review of the available scientific data and other relevant information on carboxin, the Agency has made the following determinations:

1. Pesticide products containing this sole active ingredient may be registered, subject to the terms and conditions specified in this Standard.
2. None of the risk criteria for determining unreasonable effects has been met or exceeded for this active ingredient.
3. The registrant must develop or agree to develop additional data, specified in Chapter III, to maintain the existing registration or to permit new registrations.
4. Tolerances for registered uses on barley, wheat, oats, corn, cotton, and peanuts are supported by the residue and toxicology data submitted for seed treatment uses. Tolerances are not supported for the combined foliar and soil application on peanuts (SLN use).

D. REGULATORY RATIONALE

Exposure via drift of the chemical is possible because the granular formulation is broadcast aerially for the combined foliar and soil treatment to peanut plants. Although the flowable concentrate formulation is applied by overhead irrigation or sprayed as a ground application, exposure is expected to be minimal due to the dilutions used and the use directions that state that the spray is to be directed at the crown of the plant, thereby minimizing possible drift in this use pattern.

For the predominant use pattern, dermal exposure from handling treated seed is expected to be low because most seeds are mechanically planted.

Potential exposure to wildlife exists through the ingestion of treated seeds and granules. Mechanically planted seeds may be left uncovered or partially uncovered at the end of rows, and therefore wildlife, especially birds, could ingest treated seeds.

Because there are no registered public health related uses, the Agency has waived the need for efficacy data to support the uses of carboxin.

The Agency has concluded that it should continue the registration for this chemical for the following reasons:

1. No significant adverse effects data of regulatory concern have been uncovered in the review of the studies which have been received, either in studies sufficient to meet data requirements or in those which fail to meet data requirements but which provide some level of qualitative information.
2. No significant potential risk associated with the use of carboxin is indicated from the accident data as reported on this chemical.
3. In accordance with FIFRA, the Agency's policy is not to cancel or to withhold registration merely for the lack of data. (See sections 3(c)(2)(B) and 3(c)(7) of FIFRA). Rather, publication of the Standard provides a mechanism for identifying data needs, and registration under the Standard allows for upgrading of labels during the period in which the required data are being generated. When these data are received, they will be reviewed by the Agency. The Agency will then determine whether these data will affect the registration of this chemical.

E. CRITERIA FOR REGISTRATION UNDER THE STANDARD

To be subject to this Standard, products must meet the following conditions:

- contain carboxin as the sole active ingredient; and
- bear required labeling; and
- conform to the acute toxicity limits, product composition and use pattern requirements stated in Section F, below.

The applicant for registration or reregistration of products subject to this Standard must comply with all terms and conditions described in this Standard including a commitment to fill data gaps on a time schedule specified by the Agency and, when applicable, offering to pay compensation to the extent required by 3(c)(1)(D) of the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, 7 U.S.C. 136(c)(1)(D). As discussed in chapter I, applicants for registration under this Standard must contact the Registration Division for specific instructions, including updated information contained in the guidance package on data requirements, and companies whose data must be cited and to whom compensation must be offered.

F. ACCEPTABLE RANGES AND LIMITS

1. Manufacturing-Use Products

a. Product Composition Standard

Currently the Agency has minimal information that indicates ranges and limits for the product composition of manufacturing-use products

containing carboxin. To be covered under this Standard, registrants of manufacturing-use products containing carboxin must certify ranges and limits for both active and inert ingredients, and impurities.

b. Acute Toxicity Limits

The Agency will consider registration of manufacturing-use products containing carboxin regardless of the toxicity category, provided that the labeling of such products bears appropriate precautionary statements.

c. Use Patterns and Application Methods

To be registered under this standard, manufacturing-use products containing carboxin must be labeled to allow for formulation only into end-use products which are intended for use as:

- 1) application to seeds of barley, corn, wheat, oats, cotton, or peanuts, and/or,
- 2) combined foliar and soil application to peanuts.

2. End-Use Products

a. Product Composition Standard

Currently the Agency has minimal information that indicates ranges and limits for the product composition of end-use products containing carboxin. To be covered under this Standard, registrants of end-use products containing carboxin must certify ranges and limits for both active and inert ingredients.

b. Acute Toxicity Limits

The Agency will consider registration of end-use products containing carboxin for treating seeds and combined foliar and soil application under a general-use classification, provided that they bear appropriate precautionary labeling.

c. Use Pattern and Application Method Limits

To be registered under this standard, end-use products containing carboxin must be labeled as a systemic fungicide for one or more of the following uses: application to seeds of barley, oats, wheat, corn, cotton, and peanuts. The data on combined foliar and soil application on peanuts do not adequately support registration for these uses under section 3 of FIFRA.

The full range of acceptable use patterns, maximum application rates, preharvest intervals, application methods, and other conditions are specified in Table 1.

TABLE 1
Dosage Rates and Use Limitations

<u>SITE</u>	<u>MAXIMUM DOSAGE</u>	<u>USE LIMITATIONS</u>
Barley (grain crop)	2.25 oz. a.i./100 lb. seed	Seed treatment only
Barley (seed crop)	3.00 oz. a.i./100 lb. seed	Seed treatment only
Cotton	6.00 oz. a.i./100 lb. seed	Seed treatment only
Peanuts	4.50 oz. a.i./100 lb. seed	Seed treatment only
Oats	1.12 oz. a.i./100 lb. seed	Seed treatment only
Wheat (grain crop)	2.25 oz. a.i./100 lb. seed	Seed treatment only
Wheat (seed crop)	3.00 oz. a.i./100 lb. seed	Seed treatment only
Peanuts	1.0 lb. a.i./acre	Granular combined foliar and soil application use only. ^a Preharvest interval - 20 days. ^b
Peanuts	16 oz. a.i./acre	Ground spray only. ^a Preharvest interval - 60 days. ^c
Peanuts	10.9 oz.a.i./acre	Overhead irrigation only. ^a Preharvest interval - 60 days. ^c

a/ Combined foliar and soil application to peanuts can be registered under this Standard provided (1) adequate residue data are developed to show that the present tolerance is adequate to cover the proposed use, or (2) 40 CFR 180.301 is revised to accommodate the residues occurring from the use on peanuts, peanut hulls, and peanut hay.

b/ Currently registered only as a Special Local Need (24(c)) registration in Georgia, North Carolina, Oklahoma, and Texas.

c/ Currently registered only as a SLN registration in Alabama, Georgia, North Carolina, Oklahoma, South Carolina, and Texas.

G. REQUIRED LABELING

All manufacturing-use and end-use products containing carboxin must bear labeling as specified in 40 CFR 162.10. The guidance package for this Standard contains specific information regarding label requirements. In addition, the following specific labeling requirements may apply:

1. Manufacturing-Use Products

a. Use Pattern Statements

All manufacturing-use products containing carboxin must list on the label the intended end-uses of formulated products produced from manufacturing-use products. These statements are necessary to identify the applicability of certain of the data requirements which are necessary only in support of manufacturing-use products intended for formulation into products labeled for combined foliar and soil application. These products must bear one or both of the following statements on the label:

1. "For formulation into products intended for fungicide treatment of seeds.
2. "For formulation into products intended for combined foliar and soil fungicide application."

b. Precautionary Statements

There are no unique precautionary statements which must appear on labeling of manufacturing-use products containing carboxin. The guidance package provides an updated list of all required precautionary statements typical for this type of product. The Agency may, after review of the data required under this Standard, impose additional label requirements.

2. End-Use Products

a. Precautionary Statements

For all seed treatment uses:

"Do not use treated seed for food, feed, or oil purposes."

For cotton:

"Do not graze or feed livestock on hay grown from treated seed."

For peanuts (combined foliar and soil application only):

"Do not hog down treated peanut fields prior to harvest." or

"Do not graze or feed livestock on treated areas."

For barley, corn, oats, and wheat:

"Do not graze or feed livestock on treated areas for six weeks after planting."

The guidance package provides an updated list of all required precautionary statements typical for this type of product. The Agency may, after review of the data required under this Standard, impose additional label requirements.

H. TOLERANCE REASSESSMENT

Tolerances reflecting seed treatment use have been established for combined residues of the fungicide carboxin and its metabolite 5,6-dihydro-3-carboxanilide-2-methyl-1,4-oxathiin-4-oxide (calculated as carboxin) in or on raw agricultural commodities as follows: 0.5 ppm in or on forages of barley, oats and wheat; 0.2 ppm (negligible residue) in or on cottonseed, peanuts, peanut hay and peanut hulls; 0.2 ppm in or on barley, (grain and straw), corn (grain, fodder and forage), fresh corn including sweet corn (kernels plus cob with husk removed), oats (seed and straw), rice and rice straw, sorghum (grain, fodder and forage), soybeans, and wheat (grain and straw): 0.1 ppm in meat, fat and meat by-products of cattle, goats, hogs, horses, poultry and sheep; 0.01 ppm in eggs; and 0.01 ppm in milk (40 CFR 180.301). Codex Maximum Residue Limits (MRL) have not been established for carboxin. There are no Canadian or Mexican tolerances.

Data summarized in this Standard support the established tolerances on corn, cottonseed, peanuts, sorghum, barley, oats, wheat, meat and meat by-products, and milk. Established tolerances on soybeans and rice are supported by residue data obtained with multiple-active ingredient formulations. The establishment of tolerances in poultry, meat, meat by-products and fat, and eggs are based on the conclusion that it is not possible to establish with certainty whether finite residues will be incurred, but there is no reasonable expectation of finite residues. In part, this conclusion is supported by the absence of observed bioaccumulation in a ruminant feeding study.

According to available data, the seed treatment use of carboxin falls under category (2) of 40 CFR 180.6(a), i.e. while it is not possible to establish with certainty whether finite residues in meat, milk, poultry, and eggs will result from the feeding of commodities grown from carboxin-treated seed, there is a reasonable expectation of finite residues based on the feeding of exaggerated levels of carboxin.

Carboxin tolerances on crops are based on residue data obtained with a gas chromatographic method that does not determine insoluble anilide complexes. However, residues in growing crops, analyzed by a colorimetric method that determines both soluble and insoluble residues, decline at a rate such that residues in mature crops would not be expected to exceed tolerance levels.

In calculating the theoretical maximum residue contribution (TMRC), the Agency assumes that the chemical is applied to all commodities on which its use is registered, that each commodity contains the maximum level allowed by the tolerance, and average 60 kg. adult eating patterns result in 100 percent of carboxin-treated foods in the adult diet of 1.5 kg./day. The TMRC of carboxin to the human diet is calculated to be 0.071 mg/day. The Agency has calculated an acceptable daily intake (ADI) value of 0.4 milligrams of carboxin per kilogram of body weight per day. This value is based on a "no-observed-effect" level (NOEL) of 200 ppm established in a two year rat feeding study, and the incorporation of a 100 fold safety factor in translating the data from animal to man.

Based on the TMRC of carboxin to the diet and the fact that the available data do not indicate an unreasonable effect in test animals, the established tolerances (for seed treatment only) are considered adequate to protect public health.

It must be emphasized that the established tolerances provide only for the seed treatment use of carboxin. Combined foliar and soil application of carboxin to peanuts which is registered in six states under Section 24(c) of FIFRA, may result in higher residues than those from seed treatment, thereby necessitating a higher tolerance. Additional data to evaluate this SLN use are required.

III. SUMMARY OF DATA REQUIREMENTS AND DATA GAPS

Applicants for registration of manufacturing-use and end-use carboxin products must cite or submit the information identified as required in the tables in this chapter. The tables applicable to end-use products indicate whether the product to be tested is the technical grade or formulation. Data generated on one formulation may be used to satisfy the data requirement for a substantially similar formulation. Information on which product specific data requirements are already met is available in the guidance package.

Before each requirement is listed the section of the Proposed Guidelines which describes the type of data and when the data are required to be submitted [43 FR, 29696 of July 10, 1978; and 43 FR, 37336 of August 22, 1978]. Justification for why the test is required is provided in the Guidelines. A discussion of why data additional to those already submitted are necessary, or why data normally required are not necessary for this chemical, are explained in footnotes to the tables. The data requirements specified are the minimum that will be required. Areas where additional data may be required as the result of tiered testing are indicated.

CARBON

A-1. Generic Data Requirements: Environmental Fate (See Chapter V)

Guidelines Citation	Name of Test	Composition	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation	Must Additional Data be Submitted Under FIFRA 3(c)(2)(B)? If So, due when?
163.62-7(b)	Hydrolysis	1/	no	-	yes, 14 months
163.62-7(c)	Photodegradation	1/	partial	00003088 ^{2/}	yes, 14 months
163.62-8(b)	Aerobic Soil Metabolism	1/	partially ^{3/}	00005540 00003225 05002176 00003041 00002935 05004996	yes, 14 months
163.62-8(c)	Anaerobic Soil Metabolism	1/	partially ^{4/}	00003226	yes, 14 months
163.62-8(f)	Microbial Metabolism:				
	(2) Effects of Microbes on Pesticides	-	partially	00005540 05004129 05005110 05006789 05003218	reserved ^{5/}
	(3) Effects of Pesticides on Microbes	-	partially	00005540 05002757 05003947 05003657 05003852	reserved ^{5/}

These data requirements are current as of August, 1981. Refer to the guidance package for updated requirements.

1/ Radiolabeled analytical grade or non radiolabeled technical material.

2/ This study is useful and fills part of the requirements i.e., it provided information on rate of photodegradation under natural and artificial lights. However it failed to identify the photo products. Thus further data are required. A study of photolysis on soil is also required.

3/ Data are required on aerobic soil metabolism of carboxin sulfoxide (preferably in silt loam, loamy sand or sandy soils). The data are insufficient to evaluate carboxin sulfoxide persistence in aerobic soil.

4/ Additional data are required to include 2 more soil types (preferably loam and silt loam soils).

5/ Requirements for the submission of these data are currently being reserved pending development of protocols by the Agency.

CARBOXIN

A-2. Generic Data Requirements: Environmental Fate (See Chapter V)

Guidelines Citation	Name of Test	Composition	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation	Must Additional Data be Submitted Under FIFRA 3(c)(2)(B)? If So, due when?
163.62-8(g)	Activated Sludge Metabolism	-	-	-	reserved ^{1/}
163.62-9(b)	Leaching	2/		00003227 00003114 00003229	no ^{3/}
163.62-9(d)	Adsorption/Desorption	2/	all	00005541 05003915	no
163.62-10(b)	Terrestrial Field Dissipation:				
	(1) Field & Vegetable Crops	Representative Formulation	partially ^{4/}	00003087	yes, 14 months
163.62-11(b)	Accumulation in Rotational Crops	5/	partially ^{6/}	00003114 00003224	yes, 14 months
163.62-11(d)	Fish Accumulation	2/	all	00005544 00005545	no
163.62-13	Disposal & Storage	-	-	-	reserved ^{1/}

These data requirements are current as of August, 1981. Refer to the guidance package for updated requirements.

1/ Requirements for the submission of these data are currently being reserved pending development of protocols by the Agency.

2/ Radiolabeled analytical grade or non radiolabeled technical material

3/ Data requirements have been met by data submitted on adsorption/desorption.

4/ Dissipation and mobility rates for carboxin formulations under actual use conditions are required.

5/ Radiolabeled analytical grade, and if radioactivity is found in the crops, followed with a field study using a typical representative formulation product.

6/ The available data satisfy the lab data requirements, however a field study is needed to determine if residues will be taken up by rotational crops.

August 1981

19

CARBOXIN

A-1. Generic Data Requirements: Toxicology (See Chapter VI)

Guidelines Citation	Name of Test	Composition	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation	Must Additional Data be Submitted Under FIFRA 3(c)(2)(B)? If So, due when?
163.81-1	Acute Oral Toxicity	Technical Grade of Active Ingredient	all	00003065	no
163.81-2	Acute Dermal Toxicity	Technical Grade of Active Ingredient	all	00003066	no
163.81-3	Acute Inhalation Toxicity	Technical Grade of Active Ingredient	partially ^{1/}	00003116	yes, 14 months
163.81-4	Primary Eye Irritation	Technical Grade of Active Ingredient	no	-	yes, 14 months
163.81-5	Primary Skin Irritation	Technical Grade of Active Ingredient	all	0003119	no
163.81-6	Dermal Sensitization	Technical Grade of Active Ingredient	no	-	yes, 14 months
163.82-1	Subchronic 21-Day Oral	Technical Grade of Active Ingredient	all	00003063 00003030	no
163.82-2	Subchronic 21-Day Dermal Toxicity	Technical Grade of Active Ingredient	all	00003216	no
163.82-4	Subchronic Inhalation Toxicity	Technical Grade of Active Ingredient	no	-	reserved ^{2/}

These data requirements are current as of August, 1981. Refer to the guidance package for updated requirements.

1/ Since the study gave neither the particle size nor the actual concentration of carboxin in the inhalation chamber, further testing is required.

2/ Will depend on results of an acute inhalation test for technical carboxin.

CARBOKIN

A-2. Generic Data Requirements: Toxicology (See Chapter VI)

Guidelines Citation	Name of Test	Composition	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation	Must Additional Data be Submitted Under FIFRA 3(c)(2)(B)? If So, due when?
163.83-1	Chronic Feeding	Technical Grade of Active Ingredient	all	00003031 00003152	no
163.83-2	Oncogenicity	Technical Grade of Active Ingredient	partially	00003031 00003152	yes ^{1/} , 38 months
163.83-3	Teratogenicity	Technical Grade of Active Ingredient	partially	00003120	yes ^{2/} , 24 months
163.83-4	Reproduction	Technical Grade of Active Ingredient	all	00003032	no
163.84-2 through -4	Mutagenicity	Technical Grade of Active Ingredient	partially	00003118	yes ^{3/} , 24 months
163.85-1	Metabolism (Identification of Metabolites)	Technical Grade of Active Ingredient	all	00002945 00002943 00002944	no

These data requirements are current as of August, 1981. Refer to the guidance package for updated requirements.

1/ An eighteen-month mouse oncogenicity study is needed to meet this requirement.

2/ A teratogenicity test is needed on a second mammalian species, i.e. in addition to the test on rats.

3/ Test choices within these categories must be accompanied with rationale.

(1) At least 1 more test for detecting gene mutations from among these types:

• Insects e.g. sex-linked recessive lethal test.

• Mammalian somatic cells in culture with and without metabolic activation.

• Mouse specific locus test.

(2) At least 3 test for detecting chromosomal aberrations (see 163.84-1(b)(2)(ii)).

(3) At least 2 tests for detecting primary DNA damage (see 163.84-1(b)(2)(iii)).

CARBOXIN

A-1. Generic Data Requirements: Residue Chemistry (See Chapter VII)

Name of Test	Composition	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation	Must Additional Data be Submitted Under FIFRA 3(c)(2)(B)? If So, due when?
Metabolism in Plants	Radiolabeled Active Ingredient	partially ^{1/}	05001172 05001304 05001302 05002793 05006363 05013368 05002886 05003664 05003673 05003663 05002177 00003044 00002941	yes ^{1/} , 14 months
Metabolism in Animals	Radiolabeled Active Ingredient	partially		yes ^{2/}
Analytical Methods	Technical Grade of Active Ingredient	partially	05002737 ^{3/} 00003058 ^{3/} 00003335 ^{3/} 00002919 00003054 00002905 00002940 ^{5/} 00002857 ^{5/} 00025467 00025468 00025483	yes ^{4/}

^{1/} There are sufficient data for seed treatment use; however, only partial data have been submitted for the combined foliar and soil use on peanuts, new data will be necessary.

^{2/} Additional studies to determine the adequacy of milk and meat tolerances may be necessary depending on the outcome of the peanut plant metabolism and residue studies.

^{3/} Suitable for obtaining residue data, but not suitable for tolerance enforcement.

^{4/} Additional studies may be necessary if new metabolites are identified by the metabolism studies.

^{5/} This method is acceptable for tolerance enforcement on meat, milk, and eggs. This same method may be acceptable for enforcement of tolerances on crops if supported by validation data that showed detection of insoluble anilide complexes.

August 1981

CARBOXIN

A-2. Generic Data Requirements: Residue Chemistry (See Chapter VII)

Name of Test	Composition	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation	Must Additional Data be Submitted Under FIFRA 3(c)(2)(B)? If So, due when?
Residue Data: Crop ³ -			00003335 05002737	
Corn	Technical Grade of Active Ingredient	all	00005852 00003356 00025483	no
Cottonseed	Technical Grade of Active Ingredient	all	00003129 00003185 00025468	no
Peanuts	Technical Grade of Active Ingredient	partially ^{1/}	00003045 00025468 00002905 00002903 00003300	reserved ^{2/}
Wheat	Technical Grade of Active Ingredient	all	00003219 00003158 00003045 00003218 00002961	no
Barley	Technical Grade of Active Ingredient	all	00003158 00025468 00003221	no
Oats	Technical Grade of Active Ingredient	all	00025468 00003220 00003158	no
Sorghum	Technical Grade of Active Ingredient	all	00003045 00003054 00025468	no

^{1/} Only partial data have been submitted for the combined foliar and soil use on peanuts, new data will be necessary.

^{2/} Studies may be necessary depending on the outcome of peanut plant metabolism and residue studies.

August 1981

CARBOKIN

A-3. Generic Data Requirements: Residue Chemistry (See Chapter VII)

Name of Test	Composition	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation	Must Additional Data be Submitted Under FIFRA 3(c)(2)(B)? If So, due when?
Residue Data: Processed Crops -				
Cotton seed	Technical Grade of Active Ingredient	all		no
oil			00002938	
meal			00002938	
Peanuts	Technical Grade of Active Ingredient	partially		reserved $\frac{1}{2}$, 14 months
meal			00003300	
oil			00003300	
Residue Data:				
milk	Technical Grade of Active Ingredient	partially $\frac{1}{2}$	00002945	reserved $\frac{1}{2}$, 14 months
meat	Technical Grade of Active Ingredient	partially $\frac{1}{2}$	00002945	reserved $\frac{1}{2}$, 14 months
poultry and eggs	Technical Grade of Active Ingredient	no $\frac{1}{2}$		reserved $\frac{1}{2}$, 14 months
Storage Stability	Technical Grade of Active Ingredient	yes	00025466 GS001201	no
<u>$\frac{1}{2}$ These tests may be required pending the outcome of the metabolism and residue studies on peanuts.</u>				

CARBOKIN

A. Generic Data Requirements: Ecological Effects (See Chapter VIII)

Guidelines Citation	Name of Test	Composition	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation	Must Additional Data be Submitted Under FIFRA 3(c)(2)(B)? If So, due when?
163.71-1	Avian Single-Dose Oral LD ₅₀	Technical Grade of Active Ingredient	no	-	yes, 14 months
163.71-2	Avian Dietary LC ₅₀	Technical Grade of Active Ingredient	partial ^{1/}	00003139	yes, 14 months
163.72-1	Fish Acute LC ₅₀	Technical Grade of Active Ingredient	no	-	yes, 14 months
163.72-2	Acute Toxicity to Aquatic Invertebrates	Technical Grade of Active Ingredient	no	-	yes, 14 months
163.122-1	Seed Germination	Technical Grade of Active Ingredient	no	-	yes ^{2/} , 24 months
163.122-1	Vegetative Vigor	Technical Grade of Active Ingredient	no	3/	yes ^{2/} , 24 months
163.122-2	Effects on Algae	Technical Grade of Active Ingredient	no	3/	yes ^{2/} , 24 months
163.122-2	Effects on Aquatic Macrophytes	Technical Grade of Active Ingredient	no	-	yes ^{2/} , 24 months

These data requirements are current as of August, 1981. Refer to the guidance package for updated requirements.

^{1/} Data are required on the avian dietary LD₅₀ for an upland game bird. The requirement has been fulfilled for the mallard duck.

^{2/} These data are required unless the manufacturing-use product is labeled to exclude manufacturing into products intended for combined foliar and soil application to peanuts.

^{3/} The Agency has evaluated a few studies, but they are insufficient to fulfill the requirements.

Carboxin

B. Product Specific Manufacturing-Use Products Data Requirements: Product Chemistry (See Chapter IV)

Guidelines Citation	Name of Test	Composition	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation	Must Additional Data be Submitted Under FIFRA 3(c)(2)(B)? If So, due when?
163.61-3	Product Identity & Disclosure of Ingredients	Each Product	yes	<u>1/</u>	no
163.61-4	Description of Manufacturing Process	Each Product	partially ^{2/}	-	yes ^{3/}
163.61-5	Discussion on Formation of Unintentional Ingredients	Each Product	no	-	yes ^{3/}
163.61-6	Declaration & Certification of Ingredients Limits	Each Product	no	-	yes ^{3/}
163.61-7	Product Analytical Methods & Data	Each Product	partially ^{4/}	00003172 00002995	yes, 14 months
163.61-8	Physical/Chemical Properties	Technical or Manufacturing-Use Product	partially	<u>1/</u>	yes ^{3/}

These data requirements are current as of August, 1981. Refer to the guidance package for updated requirements.

1/ Confidential Statements of Formula, EPA Form 8570-4.

2/ The manufacturing process is not sufficiently detailed for technical carboxin or the formulation intermediate.

3/ These requirements must be fulfilled by each applicant. Data from other applicants may not be cited. Therefore, even if the requirement has been partially or completely fulfilled for some products, no references are given. These requirements must be filled at the time of registration or reregistration.

4/ The analytical methods are of sufficient detail to satisfy the Agency's requirements. However, validation data and results of analysis on at least five typical samples of each product must be submitted

Carboxin

C. Product Specific End-Use Products Data Requirements: Product Chemistry (See Chapter IV)

Guidelines Citation	Name of Test	Composition	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation	Must Additional Data be Submitted Under FIFRA 3(c)(2)(B)? If So, due when?
163.61-3	Product Identity & Disclosure of Ingredients	Each Product	partially ^{1/}	2/	yes ^{3/}
163.61-4	Description of Manufacturing Process	Each Product	partially ^{4/}	-	yes ^{3/}
163.61-5	Discussion on Formation of Unintentional Ingredients	Each Product	partially ^{1/}	2/	yes ^{3/}
163.61-6	Declaration & Certification of Ingredients Limits	Each Product	no	-	yes ^{3/}
163.61-7	Product Analytical Methods & Data	Each Product	partially ^{1/}	2/	yes ^{3/}
163.61-8	Physical/Chemical Properties	Each Product	partially ^{1/}	2/	yes ^{3/}

These data requirements are current as of August, 1981. Refer to the guidance package for updated requirements.

1/ Only data specified in the Confidential Statements of Formula EPA Form 8570-4 will be required.

2/ Confidential Statements of Formula, EPA Form 8570-4.

3/ These requirements must be fulfilled by each applicant. Data from other applicants may not be cited. Therefore, even if the requirement has been partially or completely fulfilled for some products, no references are given. These requirements must be filled at the time of registration or reregistration.

4/ Manufacturing process for Vitavax Flowable Fungicide is the only process adequately described.

August 1981

27

CARBOKIN

C. Product Specific End-Use Products Data Requirements: Toxicology (See Chapter VI)

Guidelines Citation	Name of Test	Composition	Does EPA Have Data to Partially or Totally Satisfy this Requirement?	Bibliographic Citation	Must Additional Data be Submitted Under FIFRA 3(c)(2)(B)? If So, due when?
163.81-1	Acute Oral Toxicity	Each Formulation ^{1/} or Substantially Similar Formulation	partially	00003317 ^{2/} 00003081 ^{4/} 00003082 ^{4/} GS001205 ^{5/}	yes ^{3/} , 14 months
163.81-2	Acute Dermal Toxicity	Each Formulation ^{1/} or Substantially Similar Formulation	partially	00003314 ^{2/} 00003080 ^{4/} 00005858 ^{6/} 00005866 ^{8/} GS001203 ^{5/}	yes ^{7/} , 14 months
163.81-3	Acute Inhalation Toxicity	Each Formulation ^{1/} or Substantially Similar Formulation	partially	00003034 ^{2/} 00003083 ^{4/} GS001204 ^{5/}	yes ^{3/} , 14 months
163.81-4	Primary Eye Irritation	Each Formulation ^{1/} or Substantially Similar Formulation	partially	00003301 ^{2/} 00003035 ^{2/} 00003161 ^{4/} 00005857 ^{6/} 00003005 ^{8/} GS001202 ^{5/}	yes ^{7/} , 14 months
163.81-5	Primary Skin Irritation	Each Formulation ^{1/} or Substantially Similar Formulation	partially	00003311 ^{2/} 00003162 ^{4/} 00005856 ^{6/} 00005864 ^{8/} GS001206 ^{5/}	yes ^{7/} , 14 months

These data requirements are current as of August, 1981. Refer to the guidance package for updated requirements.

1/ See Guidance Package for requirements for each formulation or substantially similar formulation.

2/ This study is adequate for the testing of a 75% a.i. wettable powder/dust formulation.

3/ Further testing is required for a 29.52% a.i. soluble concentrate/liquid, 25% a.i. dust, and a 10% a.i. granular formulation.

4/ This study is adequate for the testing of a 34% a.i. ready-to-use formulation.

5/ This study is adequate for the testing of a 1% a.i. flowable formulation.

6/ This study is adequate for the testing of a 29.52% a.i. soluble concentrate/liquid formulation.

7/ Further testing is required for a 25% dust formulation.

8/ This study is adequate for the testing of a 10% a.i. granular formulation.

August 1981

IV. PRODUCT CHEMISTRY

- A. Chemical Identity
- B. Manufacturing Process
- C. Ingredient Limits in Carboxin Products
- D. Product Analytical Methods and Data
- E. Physical and Chemical Properties
- F. Summary of Data Gaps

A. CHEMICAL IDENTITY

Carboxin is the common name accepted by the American National Standards Institute (ANSI) for the chemical 5,6-dihydro-2-methyl-1,4-oxathiin-3-carboxanilide. Carboxin is also known by the trade name Vitavax® and by the abbreviations DMOC and DCMO. The Chemical Abstracts Registry number is 5234-68-4; the Uniroyal, Inc. internal code numbers as D-735 and F-735; and the EPA Shaughnessy number is 090201. The common name, carboxin, will be used throughout this standard in lieu of other chemical or trade names.

B. MANUFACTURING PROCESS

The specific details of the synthesis process for technical carboxin are considered trade secrets. There are two procedures, and they are detailed in Uniroyal, 1977, MRID 00003296 and Uniroyal, 1976, MRID 00003084. The manufacturing process for the formulation - Vitavax® Flowable Fungicide - has been submitted to the Agency (Uniroyal, 1977, MRID 00003231). This process is also considered a trade secret.

Manufacturing processes for the other products have not been submitted to the Agency.

C. INGREDIENT LIMITS IN CARBOXIN PRODUCTS

A commitment (certification) is required from each registrant that the ingredients and impurities in the product will be maintained within specified limits for as long as the product is offered for sale. Upper and lower limits are required for the active and intentionally added inert ingredients. Upper limits are required for the impurities. These have not been established for any carboxin product.

D. PRODUCT ANALYTICAL METHODS AND DATA

Infrared spectroscopic methods for the assay of carboxin in the technical product and in formulations have been submitted to the Agency, (Puchalsky, 1968, MRID 00003172 and Uniroyal, 1977, MRID 00002995). A titrimetric method has also been submitted for the determination of carboxin in the technical product (Uniroyal, 1960, MRID 00002978). The above-mentioned infrared spectroscopic and titrimetric methods have been published by Stone (1976, MRID 05005076).

Carboxin formulations may be analyzed for the active ingredient by an ultraviolet spectrophotometric method (Harda, 1978, MRID 05003778).

The above mentioned assay methods are sufficiently detailed to satisfy Agency requirements. However, validation data and results of analysis on at least five typical samples of each product have not been submitted. This information will need to be submitted.

E. PHYSICAL AND CHEMICAL PROPERTIES

The following information is for technical carboxin, unless otherwise mentioned.

1. Color

Off-white (Uniroyal, 1977, MRID 00005859 and Confidential Statement of Formula [CSF])

2. Odor

Described as "faint" (Uniroyal, 1977, MRID 00005859 and CSF)

3. Melting Point

Two melting ranges of technical carboxin are 91.5-92.5°C and 98-101°C, reflecting two crystalline structures. In solution the two structures revert to one. It is reported that there are no differences in biological activity between the two structures (Uniroyal, 1977, MRID 00005859 and CSF).

4. Density or Specific Gravity

<u>Product</u>	<u>Density/Specific Gravity</u>
Technical	1.70
Technical	40-45 lbs/ft ³ *
Formulation Intermediate	20-30 lbs/ft ³ *
Ready to Use (Flowable)	1.05-1.13 gm/ml
Soluble Concentrate (Liquid)	1.13-1.18 gm/ml
	*Bulk density
	(Uniroyal, 1977, MRID 00005859 and CSF)

5. Physical State

<u>Product</u>	<u>Physical State</u>
Technical	Crystalline solid
Formulation Intermediate	Solid
Wettable Powder/Dust	Powder
Ready-to-Use (Flowable)	Liquid
Soluble Concentrate	Liquid
Dust	Powder

(Uniroyal, 1977, MRID 00005859 and CSF)

6. Solubility

<u>Solvent</u>	<u>Gm Solute/100 gm solvent @ 25 C</u>
Distilled Water	0.017
Benzene	15
Dimethyl Sulfoxide	150
Acetone	60
Methanol	21
Ethanol	11

(Uniroyal, 1977, MRID 00005859)

7. Stability

Carboxin is readily inactivated by ultraviolet light and sunlight (Buchenour, 1975, MRID 05002823).

8. pH

There are two Ready-to-Use (Flowable) formulations. The pH range for one of these formulations is 7.0-9.0, the other is 6.2-8.2. The pH for the soluble concentrate formulation is 9.0-10.5 (Uniroyal, 1977, MRID 00005859 and CSF).

9. Flammability

<u>Product Type</u>	<u>Flash Point</u>	<u>Test Method</u>
Technical	203 C	C.O.C.*
Ready-to-Use (Flowable)	>93 C	Unreported
Ready-to-Use (Flowable)	102 C	Unreported
Soluble Concentrate (Liquid)	56 C	Unreported

*Cleveland Open Cup Method

(Uniroyal, 1977, MRID 00005859 and CSF)

10. Storage Stability

Three years (Uniroyal, 1977, MRID 00005859).

F. SUMMARY OF DATA GAPS

A detailed manufacturing process for the technical product, the formulation intermediate, and each end-use product except Vitavax® Flowable Fungicide; details on the formation of unintentional ingredients; certification of ingredient limits; validation data and results from analysis on each product; and various physical/chemical properties must be submitted to the Agency.

V. ENVIRONMENTAL FATE

- A. Use Summary
- B. Environmental Fate Profile
- C. Exposure Profile
- D. Summary of Data Gaps

A. USE SUMMARY

Carboxin is a systemic fungicide registered for use as a seed treatment for control of smuts and seed rot and seedling blight caused by Rhizoctonia solani. As a single active ingredient, use sites include barley, corn, cottonseed, oats, peanuts, and wheat. Carboxin is also registered for Special Local Needs (FIFRA section 24(c)) as a combined foliar and soil application on peanuts to control Southern blight (in Alabama, Georgia, North Carolina, Oklahoma, South Carolina, and Texas only).

At present six products with carboxin as the sole active ingredient are registered for use: a dust (25% a.i.); a wettable powder/dust (75% a.i.); two ready-to-use flowable liquids (34% and 17.1% a.i.); and two liquid soluble concentrates (both 29.5% a.i.). For SLN use there are two formulations registered: a 34% flowable concentrate and a 4% granular.

Current seed treatment uses and application rates, by formulation, are summarized in Table 1; and SLN combined foliar and soil application rates are summarized in Table 2, on the following pages.

Table 1
Seed Treatment Application Rates

Formulation*	RATE RANGE (oz. product/100 lb seeds)			
	WP/D (75%)	RTU (34% & 17.1%)	D (25%)	SC/L (29.52%)
<u>Crop</u>				
Barley	2-3 ^a /	2-3 ^b /		
Corn		2-4 ^b /	4-6	
Cottonseed	4-8	16 ^c /		20 ^d /
Oats	1-2	2-3 ^b /		
Peanuts	2-6			
Wheat	2-3 ^a /	2-3 ^b /		

*WP/D- Wettable Powder/Dust

RTU- Ready-to-Use

D- Dust

SC/L- Soluble Concentrate/Liquid

^a/4 oz./100 lbs. for seed production purposes only

^b/34% Ready to Use Formulation

^c/17.1% Ready to Use Formulation

^d/Limited by registrant to seed treatment by
professional applicators only

Table 2
Foliar Application Rates on Peanuts

Formulation*	FC (34%)	G (4%)
State		
Alabama	3 pints/acre ^{a/}	
Georgia	3 pints/acre ^{a/}	25 lbs./acre ^{d/}
North Carolina	3 pints/acre ^{b/}	25 lbs./acre ^{d/}
Oklahoma	3 pints/acre ^{a/}	25 lbs./acre ^{d/}
	2 pints/acre ^{c/}	
South Carolina	3 pints/acre ^{a/}	
Texas	3 pints/acre ^{a/}	25 lbs./acre ^{d/}
	2 pints/acre ^{c/}	

* FC- Flowable Concentrate

G- Granular

^{a/}Mixed with 20 gallons of water and applied by ground spray

^{b/}Mixed with 30 gallons of water and applied by ground spray

^{c/}Applied by overhead irrigation only

^{d/}Applied by air or ground broadcast

B. ENVIRONMENTAL FATE PROFILE

The available data are insufficient to completely assess the environmental fate of carboxin.

In aqueous solution (under UV light and in the dark) carboxin was oxidized to carboxin sulfoxide, carboxin sulfone, and two unidentified compounds (A and B). The photolytic half-life was less than four hours. After seven days of irradiation in aqueous solution, 40 and 9 percent of the exposed ^{14}C carboxin was present as the unidentified compounds A and B, respectively. Fourteen days later, 57 percent of the radioactivity was present as compound A and 19 percent as compound B. Formation of compound A was accelerated in a 2 percent acetone-water solution (Smilo, 1977, MRID 00003088).

Carboxin was degraded in aerobic soil. The half-life of carboxin was less than one day in loamy sand and silt loam soil, and less than three days in sandy soil, with more than 95 percent of the applied carboxin being degraded within seven days. The major degradation product was carboxin sulfoxide, which represented 31-54 percent of the applied radioactivity at seven days after treatment. Carboxin sulfoxide residue levels declined to 28-49 percent and 19 percent of the applied radioactivity at 30 and 154 days after treatment, respectively. Several minor degradation products were also formed (carboxin sulfone, p-hydroxy carboxin sulfoxide, p-hydroxy carboxin, and $^{14}\text{CO}_2$). Carboxin was degraded in sterile soil but at a much slower rate than in nonsterile soil (46-72 percent degraded in seven days). This indicated that the soil metabolism of carboxin under aerobic conditions was both a microbial and chemical process. Carboxin sulfoxide and carboxin sulfone were stable under anaerobic conditions (Chin et al., 1972, MRID 00002935; Chin et al., 1969, MRID 00003041; Chin et al., 1970, MRID 05002176; Chin et al., 1970, MRID 05004996; Dzialo and Lacadie, 1978, MRID 00003225; Dzialo et al., 1978, MRID 00003226; and Spare, 1979, MRID 00005540).

Carboxin was oxidized to carboxin sulfoxide and carboxin sulfone by flavin enzymes found in fungal mitochondria. The fungus Ustilago maydis and isolated mitochondria from Trametes versicolor and Aspergillus niger were capable of oxidizing carboxin to carboxin sulfoxide and carboxin sulfone. This reaction was accelerated under illuminated conditions. A Pseudomonas species oxidized carboxin to carboxin sulfoxide in three days and to carboxin sulfone in seven days (Balasubramanya and Patil, 1976, MRID 05006289). Carboxin sulfone was hydrolyzed to form 5,6-dihydro-2-methyl-1,4-oxathiin-3-carboxylic acid-4,4-dioxide and aniline (Lyr et al., 1974, MRID 05003852). Basillus cereus oxidized carboxin to carboxin sulfoxide and carboxin sulfone in Nile River water containing added sludge (El-Dib and Aly, 1976, MRID 05003218; Bachofer et al., 1973, MRID 05005110; and Michail et al., 1975, MRID 05004129).

The effect of carboxin on microorganisms is varied. At two ppm, carboxin had a slight inhibitory effect on fungal populations for 21 days and inhibited starch degradation (33-55 percent) for three days. A severe inhibitory effect on nitrification (75-85 percent) occurred from 7 to 14 days after treatment with carboxin at two ppm. This inhibition may have been due to carboxin sulfoxide

since more than 95 percent of the carboxin was degraded within seven days (Spare, 1979, MRID 00005540). Carboxin at 50 ppm inhibited dehydrogenase activity in fungi 30 percent (Lyr et al., 1974, MRID 05003852).

In the laboratory, carboxin inhibited two strains of nitrogen-fixing bacteria in nutrient agar. When applied to inoculated seed at 0.1-0.2 ppm (approximate actual application rate), carboxin inhibited nodulation of soybeans inoculated with *Rhizobium japonicum* by up to 83 percent and completely inhibited nodulation of *Vigna anguiculata* inoculated with rhizobia strains CB 756 and XS 30 however, treated uninoculated seeds planted in soil containing rhizobia strains CB 756 and XS 30 produced nodules (Curley and Burton, 1975, MRID 05003947). *Azotobacter chroococcum* and *Rhizobium trifolii* in nutrient agar were not inhibited by carboxin at two and ten ppm, respectively. Although carboxin sulfone at 25-150 ppm decreased nitrogen fixation by *Rhizobium trifolii* 20-25 percent, it is doubtful that carboxin sulfone will be present in concentrations greater than 0.05 ppm as a result of the application of carboxin. Of the 23 common soil bacteria and fungi exposed to carboxin, only four experienced growth reductions of 20 percent or more at the lowest concentration of 2.5 ppm (Spare, 1979, MRID 00005540; El-Dib and Aly, 1976, MRID 05003218; Fisher, 1976, MRID 05002757; and Kritzman et al., 1977, MRID 05002989).

Both carboxin and carboxin sulfoxide were mobile in soil, with about half of the applied ^{14}C carboxin leaching through the twelve inch column of clay loam soil (Lacadie et al., 1978, MRID 00003227 and Dannals et al., 1976, MRID 00003114). Lacadie et al. (1978, MRID 00003229) found that radiolabeled carboxin aged in sandy soil was mobile. Approximately 17 percent of the applied radioactivity was recovered in the leachate from a twelve inch soil column. One-third of the radioactivity was in the top three inches and one-fourth was in the three to six inch segment of the soil column. Carboxin, carboxin sulfoxide and carboxin sulfone were identified in the leachate and accounted for 0.3, 3.3 and 0.5 percent respectively of the applied radioactivity.

Smyser (1979, MRID 00005541) found that there is a low potential for carboxin adsorption to a sandy loam soil (Freundlich adsorption coefficient was $K=0.78$). The calculated desorption coefficient was $K=1.10$.

A simulated field dissipation study was conducted by Cardona et al. (1976, MRID 00003087). One month after a sandy loam soil was treated with radiolabeled carboxin (1 lb./acre) only 4 percent of the applied carboxin had not degraded. The major degradation products were carboxin sulfoxide (31-33 percent) and an unidentified compound II (6-18 percent). Two months after treatment the carboxin could not be detected and only 4 percent of the sulfoxide and 2-3 percent of compound II remained. After one year, approximately 75-80 percent of the radioactivity remaining was found in the top six inches. Leaching to at least 11 inches was indicated by the detection of radioactivity at that depth. The detection of the compound, its metabolites, or degradation products at a depth of 11 inches one year after application indicated that carboxin aged residues were persistent and mobile.

Dannals et al. (1976, MRID 00003114) found ^{14}C residues in wheat (seed), beets (top and root), and lettuce planted in a sandy loam soil four months after treated with ^{14}C -carboxin at 1 lb. a.i./acre. The concentration of oxathiin-labeled residues present in those crops were 1.5-60 times higher than the concentration of aniline-labeled residues.

A field study by Uniroyal Chemical (1978, MRID 00003224) showed that carboxin residues were not taken up in turnip roots planted 59 days after treatment with carboxin (Vitavax® 3F). Carboxin residues were less than 0.2 ppm, the sensitivity of the method used, in turnip greens and rye seed planted after treatment. However, the analytical method was not sensitive enough to determine conclusively that residues less than 0.2 ppm were not taken up by rotational crops.

Bluegill sunfish exposed to carboxin at a constant concentration of 0.012 ppm for 30 days accumulated the compound at 0.50 ppm with a bioaccumulation factor of 45. A plateau was not reached during the 30-day exposure period, thus indicating that there is a potential for carboxin residues to accumulate in sunfish. Residue levels were increasing at the end of the exposure period, indicating that continued uptake of ^{14}C residues could occur because carboxin oxidative products (carboxin sulfoxide and carboxin sulfone) are stable and persistent in water and probably would be available for uptake (accumulation) beyond 30 days. After a 14-day depuration period, 23 percent (0.11 ppm) of the maximum ^{14}C residue levels remained in the sunfish (Kuc and Doebbler, 1979, MRID 00005544). Catfish exposed to carboxin and its aged residues at 0.05 ppm accumulated 0.36 ppm with a bioaccumulation factor of 5 after exposure for 22 days. The ^{14}C residue levels in the whole fish tissue had declined by approximately 15 percent (to 0.31 ppm) at 30 days. After a 14-day depuration period, 32 percent (0.12 ppm) of the maximum ^{14}C residue levels in catfish remained (Kuc and Doebbler, 1979, MRID 00005545). Since 40-80 percent of the residues in the bluegill sunfish and over 50 percent of the residues in the catfish were unextractable, the identity of the unextractable compounds cannot be established.

In summary, carboxin is rapidly degraded to carboxin sulfoxide and carboxin sulfone (greater than 95 percent within seven days) by microbial and chemical processes in aerobic soil. Based on the mobility data available, carboxin was not tightly adsorbed to sandy loam soil and was mobile in soil columns under rapid leaching conditions, indicating a potential to contaminate ground water. Mobility of carboxin aged in soil was mitigated by its degradation to carboxin sulfoxide and carboxin sulfone. However, carboxin sulfoxide and small amounts of carboxin sulfone were mobile in soil columns, indicating their potential to reach ground water.

Carboxin severely yet briefly inhibits starch degradation and when used as a seed treatment on inoculated seeds severely inhibits formation of nodules by some strains of nitrogen-fixing bacteria. Therefore, the use of carboxin on inoculated legumes (currently used on peanuts) may have an effect on symbiotic nitrogen fixation and therefore on the amounts of nitrogen available to peanuts.

Carboxin residue levels of 0.50 and 0.36 ppm accumulated in bluegill sunfish and catfish. Carboxin sulfoxide was the major carboxin metabolite in bluegills, but the carboxin residues in catfish were not identified.

Carboxin sulfoxide was the major degradation product of carboxin in soil and water. Carboxin sulfoxide was present in aerobic soil for at least five months and was stable in anaerobic soil. Carboxin was photodegraded to carboxin sulfoxide and two unidentified compounds. Carboxin sulfoxide may inhibit nitrification in soil.

Carboxin sulfone (the fungicide oxycarboxin) is a minor product of carboxin degradation. It will not be present in soil in a sufficient concentration to result in significant exposure.

C. EXPOSURE PROFILE

1. Seed Treatment Formulations Only

Exposure from this use pattern to humans, livestock, and wildlife via spray drift is unlikely because the chemical is not applied aerially. Carboxin and its residues have been shown to be mobile in soil indicating a potential for groundwater contaminate. Although the mobility of carboxin aged in soil was mitigated by its degradation to the sulfoxide and sulfone, these degradates were also mobile in soil. However, since single active ingredient formulations of carboxin are used primarily as seed treatments, the use pattern minimizes the potential exposure of humans and domestic animals to carboxin and its residues via groundwater contamination. Carboxin residues indicated a potential to accumulate in bluegill sunfish and catfish. However, as formulations containing the single active ingredient carboxin are used primarily in seed treatments, this potential hazard is minimized. Potential exposure of humans by ingestion of carboxin residues in rotated crops is also minimized by this use pattern.

Potential exposure of wildlife exists through the ingestion of treated seeds. Mechanically planted seeds may be left uncovered or partially covered at the end of rows, and therefore wildlife, especially birds, could ingest treated seeds. Data necessary to estimate the nature and extent of such exposure are unavailable.

The greatest potential for human exposure exists during seed treatment, most of which is done by seed processors or seed companies. Respiratory exposures may be especially high from the use of dust formulations, and commercial applicators are switching to the flowable concentrate to reduce such exposure. Dermal exposure from handling treated seed is expected to be low since most seeds are mechanically planted.

2. Special Local Need Registrations

a. Flowable Concentrate

Exposure to humans, livestock and wildlife via spray drift is unlikely because the product is not applied aerially. Exposure by spray drift as a result of ground application is not expected because of the use directions which state that the spray solution should be directed to the crown of the plant and pegging zone as a course spray. Direct exposure to humans by overhead irrigation of plants is expected to be minimal due to the dilution directions involved in its application. However, dermal, ocular and oral exposure to humans may occur from drinking and washing of face and hands with irrigation water. Wildlife and domestic animals may potentially be exposed by drinking the irrigation water. Carboxin and its residues have been shown to be mobile in soil indicating a potential to runoff and/or contaminate ground water. The data necessary to develop a quantitative assessment of carboxin and its residues to contaminate ground water are not available. Carboxin residues indicated a potential to accumulate in bluegills and catfish. The Agency does not currently have data to develop a quantitative assessment of fish exposure to residues of carboxin in adjacent ponds as a result of runoff. Potential exposure of humans to carboxin residues by ingestion of rotated crops is increased by the increased dosage and the number of applications involved in the flowable concentrate use on peanuts. The Agency does not currently have data to estimate the nature and extent of such exposure.

b. Granular Formulation

Exposure is possible from the drift of dust particles from aerial applications of granulars. However, little or no information is presently available on the aerial application of granular fungicides. Potential exposure of humans to carboxin and its degradation products by ingestion of rotated crops is increased by the increased dosage and number of applications involved in the granular product use on peanuts, since the granular product falls predominantly on the ground with little intercepted by peanut foliage. Because of increased levels of residues, the preharvest interval has been increased to 60 days. Exposure by mobility and accumulation in fish is similar to the flowable concentrate, but the exposure is increased because of the use pattern for the granular product.

D. SUMMARY OF DATA GAPS

A number of the guideline requirements have been partially fulfilled by the data submitted. However, data are still needed to adequately assess the environmental fate of carboxin. The specific deficiencies can be found in the Data Requirements Charts in Chapter III. The data gaps are: hydrolysis, photodegradation, aerobic and anaerobic soil metabolism, terrestrial field dissipation, and accumulation in rotational crops.

VI. TOXICOLOGY

- A. Toxicology Profile
- B. Human and Domestic Animal Hazard Assessment
- C. Summary of Data Gaps

A. TOXICOLOGY PROFILE

1. Technical Carboxin

a. Acute Effects

A limited amount of information was available to assess the acute oral toxicity of technical carboxin. The oral LD₅₀ in rats was 3.82 ± 0.35 g/kg which is sufficient to assign technical carboxin to Toxicity Category III (Carson, 1965, MRID 00003065).

An acute dermal toxicity test was conducted in rabbits (Carson, 1965, MRID 00003066). The data show that when applied as a 50% aqueous slurry, technical carboxin causes no mortality at levels of 8 g/kg, which is sufficient to assign it to Toxicity Category III.

Babish (1977, MRID 00003116) reported that exposure for one hour to a concentration of 20 ml/l of technical carboxin did not cause mortality in rats in an acute inhalation toxicity test. However, due to inappropriate testing protocols, this study must be repeated.

In a primary dermal irritation study conducted on rabbits, 0.5 gm of technical carboxin was applied to the skin and did not cause irritation (Babish, 1977, MRID 00003119). This study indicates that technical carboxin is not a potential skin irritant and may be assigned to Toxicity Category IV.

b. Subchronic Effects

Sufficient data were available to assess the subchronic dietary and dermal effect of technical carboxin. In the rat 90-day feeding study the No Observable Effect Level (NOEL) was 200 ppm of carboxin in the diet. At the 600 ppm level, the effects noted were degenerative renal changes (Ozer, 1966, MRID 00003063). In the two-year dog feeding study the NOEL was 600 ppm (Holsing, 1969, MRID 00003030). In addition, in a 21-day dermal study, rabbits treated with 3.0 g/kg did not produce any treatment related effects (Holsing, 1968, MRID 00003216). No data were available to assess the subchronic inhalation toxicity of technical carboxin. However, the requirements of these data is reserved pending the final results of the acute inhalation study.

c. Chronic Effects

In a screening study, groups of rats were fed for two years diets containing 0, 300, 1,000 or 3,000 ppm of technical carboxin (Holsing, 1969, MRID 00003152). A NOEL was not established in this study because the dose levels were too high. However, in a subsequent chronic feeding study over a two year period,

rats were fed diets containing 0, 100, 200, or 600 ppm of technical carboxin (Holsing, 1969, MRID 00003031). This study indicated that the NOEL is 200 ppm. At the 600 ppm level the effects were poor survival and weight gain depression. These data are sufficient to satisfy the requirement for chronic feeding.

No evidence of oncogenicity was indicated in either of the two-year feeding studies (Holsing, 1969, MRID 00003031 and Holsing, 1969, MRID 00003152). However, oncongenic testing in a second species (preferably the mouse) is required.

A teratology study was conducted in rats by Knickerbocker (1977, MRID 00003120). Technical carboxin at doses as high as 4.0 mg/kg/day did not produce any maternal toxicity. However, teratogenicity testing is still required in a second species.

In a three generation reproduction study, rats were fed diets containing 100, 200, and 600 ppm technical carboxin (Holsing, 1968, MRID 00003032). There were no treatment related effects on reproductive performance. The NOEL was established at 200 ppm.

d. Mutagenicity

Mutagenicity testing is incomplete. In a study by Brusick (1977, MRID 00003118) two different types of tests for detecting gene mutations were reported. Carboxin was tested in the Salmonella typhimurium reversion assay with and without a rat liver microsome metabolic activation system. No increases in reversion frequency were detected in either assay. Carboxin was also tested in a mutation assay system using Saccharomyces cerevisiae, but procedures in this assay were inadequately described. Additional mutagenicity testing is required.

e. Metabolism

Kennedy (1971, MRID 00002943) reported on the distribution and excretion of ¹⁴C carboxin in rats. Between 88 and 99 percent of the compound was recovered, most within 24 hours of treatment. The urine contained 42-89 percent of the administered ¹⁴C and the feces contained 10-45 percent. The remainder was found in the animal's tissues and organs.

In a companion study, Kennedy (1971, MRID 00002944) characterized the ¹⁴C activity in urine and feces using thin layer chromatography and liquid scintillation. Carboxin sulfoxide accounted for 27-45 percent of the ¹⁴C in the 24 hour sample and 47-56 percent in the 72 hour sample. Three additional urinary components were detected but not identified, and the parent compound was not ¹⁴C present. The major component found in the feces (19-36 percent of fecal ¹⁴C) was tentatively identified as carboxin sulfone.

These studies collectively provide sufficient information about the metabolism of carboxin in animals.

2. End-Use Carboxin

Acute toxicity (oral, dermal, and inhalational) and irritation testing (eye and dermal) is required for each formulation or substantially similar formulation.

a. Acute Effects- Oral

One test (Matthews, 1970, MRID 00003317) is available on the acute oral LD₅₀ of a 75% wettable powder/dust formulation. The LD₅₀ value for this formulation in albino rats is greater than 2 g/kg. This is sufficient information to assign this formulation to Toxicity Category III.

Two acute oral toxicity studies were conducted that together show that the LD₅₀ for the 34% ready-to-use liquid formulation in rats is greater than 5 ml/kg, or 5 g/kg (Babish, 1977, MRID 00003081 and Babish, 1977, MRID 00003082). This is sufficient information to place this formulation in Toxicity Category IV.

The acute oral LD₅₀ of the 17% flowable formulation was found to be greater than 5 ml/kg in rats (Babish, 1977, MRID GS001205). This formulation falls into Toxicity Category IV.

b. Acute Effects-Dermal

A dermal toxicity study was conducted on rabbits using a 75% wettable powder/dust formulation (Matthews, 1970, MRID 00003314). No deaths and no signs of toxicity were observed at doses up to 10 g/kg. This is sufficient information to assign this formulation to Toxicity Category III.

The acute dermal LD₅₀ value of a 34% ready-to-use formulation was found to be greater than 20 ml/kg or 20 g/kg in albino rabbits (Babish, 1977, MRID 00003080). This is sufficient information to assign this formulation to Toxicity Category IV.

The acute dermal LD₅₀ value of a 29.52% soluble concentrate/liquid formulation was found to be greater than 5 g/kg for albino rabbits (Stevens, 1979, MRID 00005858). This information is sufficient to assign this formulation to Toxicity Category III.

The acute dermal LD₅₀ of a 10% granular formulation is greater than 20 g/kg in albino rabbits (Babish, 1977, MRID 00005866). No deaths occurred in the 14 day observation period. Decreased activity and anorexia were the only toxic signs. This is sufficient information to assign this formulation to Toxicity Category IV.

The acute dermal LD₅₀ of a 17% flowable formulation is greater than 20 g/kg in rabbits (Babish, 1977, MRID GS001203). This test substance falls within Toxicity Category IV.

c. Acute Effects- Inhalation

Binnes et al. (1969, MRID 00003034) found that the LC_{50} value of a 75% wettable powder/dust formulation is greater than 5 mg/l for rats. No animals died, but test animals displayed a nasal discharge during exposure. This formulation can be assigned a Toxicity Category III.

Rats were exposed to a concentration of 20 mg/l of a 34% ready-to-use formulation. No mortality was produced, but ataxia was observed (Babish, 1977, MRID 00003083). These data indicate that this formulation can be assigned to Toxicity Category IV.

The inhalation LC_{50} of a 17% flowable formulation is greater than 20 ml/l in rats (Babish, 1977, MRID GS001204). This test substance falls within Toxicity Category IV.

d. Acute Effects- Eye Irritation

Two studies were conducted using a 75% wettable powder/dust formulation. Holsing (1968, MRID 00003035) placed 100 mg of the formulation in the eyes of rabbits and observed marked and persistent conjunctival effects, iris irritation, and corneal opacity. In general, the irritation and corneal damage were still present after 14 days. Bailey (1976, MRID 00003301), also placed 100 mg of the 75% wettable powder formulation into one eye of each rabbit and noted corneal opacity within 72 hours, and damage to the iris, although neither of these effects persisted for seven days. Conjunctival effects occurred and persisted for seven days. This information is sufficient to assign this formulation to Toxicity Category I, however, if the registrant wishes to repeat an eye irritation study for the required 21 days, the Toxicity Category may be lowered.

Testing a 34% ready-to-use formulation for eye irritation, Babish (1976, MRID 00003161) found no irritation to the conjunctiva, iris or cornea of the eyes of the test rabbits. This formulation can be assigned to Toxicity Category IV.

Stevens (1979, MRID 00005857) tested a 29.52% soluble concentrate/liquid formulation for eye irritation. Results show transient conjunctival effects, however no iris or cornea irritation was observed. These observations are sufficient to assign this formulation to Toxicity Category IV.

A 10% granular formulation was instilled in one eye (unwashed) of each of six rabbits (Bailey, 1976, MRID 00003005). This treatment did not produce corneal damage, but did cause slight conjunctival effects, including redness, swelling, and discharge. These effects started within 72 hours in all six rabbits and persisted in four rabbits for seven days. These results are sufficient to place this formulation in Toxicity Category III, however, if the registrant wishes to repeat an eye irritation study for the required 21 days, the Toxicity Category may be lowered.

A 17% flowable formulation was a mild to moderate eye irritant to rabbits which produced corneal opacity persisting over 72 hours, but less than seven days

(Babish, 1977, MRID GS001202). The average Draize score was 0.66, which is sufficient to place this formulation into Toxicity Category III.

e. Acute Effects- Dermal Irritation

No irritation was observed at 24 and 72 hours after a 75% wettable powder/dust formulation was applied to the skin of rabbits (Matthews, 1970, MRID 00003311). These results are sufficient to assign this formulation to Toxicity Category IV.

Moderate irritation to the intact and abraded skin of albino rabbits was observed at 24 and 72 hours after application of a 34% ready-to-use formulation (Babish, 1976, MRID 00003162). Erythema and edema were also noted. These results were sufficient to assign this formulation to Toxicity Category IV.

The 29.52% soluble concentrate/liquid formulation was severely irritating when applied to both intact and abraded skin of albino rabbits. Moderate to severe erythema and edema persisted throughout the 72 hour observation period (Stevens, 1979, MRID 00005856). These results were sufficient to assign this formulation to Toxicity Category III.

No irritation was observed at 24 and 72 hours after a 10% granular formulation was applied to the skin of rabbits (Babish, 1977, MRID 00005864). These results are sufficient to assign this formulation to Toxicity Category IV.

A 17% flowable formulation produced edema for 72 hours when tested on rabbits (Babish, 1977, MRID GS001206). The average Draize score was 0.50, indicating that this formulation is not classified as a skin irritant and can be placed into Toxicity Category IV.

B. HUMAN AND DOMESTIC ANIMAL HAZARD ASSESSMENT

1. Technical

The information available to assess potential hazard as a result of chronic exposure is incomplete (see the Toxicity Profile for details). However, the available data suggest carboxin poses little or no discernible risk in regards to oncogenic, teratogenic and reproductive effects to man or domestic animals. In addition, the data on acute oral, acute dermal and primary dermal irritation indicate the products containing carboxin are in Toxicity Category III. No data were available to assess the acute inhalation, primary eye irritation and dermal sensitization hazard.

2. Wettable Powder/Dust (WP/D)

The 75% WP/D formulation had low acute oral, acute dermal, acute inhalation and primary skin irritation potentials. However, the overriding toxic effect is the primary eye irritation category. Two irritation studies show that this formulation is a moderate to severe eye irritant and appropriate labelling and caution (Category I) is necessary.

3. Ready-to-Use (RTU)

The 34% and 17% RTU formulations, according to the acute oral, acute dermal, acute inhalation, primary eye, and primary dermal irritation, are in Toxicity Category IV (low hazard potential).

4. Soluble Concentrate/Liquid (SC/L)

The 29.5% SC/L formulation, according to the acute dermal, primary eye irritation, and dermal irritation, is in Toxicity Category III. This is due to the moderate dermal irritation effects.

5. Granular

The 10% granular formulation showed low dermal toxicity and irritation, however there was moderate eye irritation. This formulation can be placed in Toxicity Category III.

C. SUMMARY OF DATA GAPS

The following tests are required for the reregistration of carboxin: acute oral toxicity (25% dust, 29.5% soluble concentrate/liquid, and 10% granular), acute dermal toxicity (25% dust), primary eye irritation (technical product and 25% dust), skin irritation (25% dust), acute inhalation toxicity (10% granular, 25% dust, 29.5% soluble concentrate/liquid), and the following tests on the technical product only: dermal sensitivity, oncogenicity, teratogenicity, and mutagenicity.

VII. RESIDUE CHEMISTRY

- A. Residue Chemistry Profile
- B. Labeling Requirements
- C. Summary of Data Gaps

A. RESIDUE CHEMISTRY PROFILE

1. Uptake and Distribution and Metabolism in Plants

Data obtained by a variety of methods show that carboxin is systemic in plants. Extracts of plants grown from carboxin-treated seed or immersed in a carboxin solution have been shown by bioassay methods to contain fungicidal residues (Verma and Vyas, 1976, MRID 05001172; Thapliyal and Sinclair, 1971, MRID 05001304; Thapliyal and Sinclair, 1970, MRID 05001302; Bolkan and Milne, 1975, MRID 05002793). Fungicidal residues persist up to 29 days in some plants (Bolkan and Milne, 1975, MRID 05002793).

The distribution of residues derived from phenyl-¹⁴C- or oxathiin-¹⁴C-carboxin in cotton, wheat, barley, soybean, and bean plants has been studied (Leroux and Gredt, 1972, MRID 05006363; Ambro-Balint, 1974, MRID 05013368; Briggs et al., 1974, MRID 05002886; Thapliyal and Sinclair, 1971, MRID 05001304; Kirk et al., MRID 1969, 05003664; Berggren and Pinckard, 1973; MRID 05003673; Snel and Edgington, 1970, MRID 05003663; Chin et al., 1970, MRID 05002177; Chin et al., 1969, MRID 00003044; Chin et al., 1972, MRID 00002941). Bioassay data show that carboxin is systemic in all species studied. The pattern of distribution of carboxin and its metabolites within plants is variable, depending upon the species examined, the length of exposure to the labeled compound. Data obtained by methods capable of detecting the total radiolabeled residue (combustion technique, radioautography) show that roots, lower stem and the earliest leaves contain the bulk of the radiolabeled chemical (Kirk et al., 1969, MRID 05003664; Berggren and Pinckard, 1973, MRID 05003673; Snel and Edgington 1970, MRID 05003663; Chin et al., 1969, MRID 00003044). Radiolabeled carboxin distribution within plants was similar when the ¹⁴C label was in the oxathiin or phenyl rings of carbon (Briggs et al., 1974, MRID 05002886; and Chin et al., 1969, MRID 00003044).

The predominant metabolite of carboxin in wheat, beans, and barley plants grown from ¹⁴C-carboxin-treated seed is the sulfoxide derivative of carboxin (Leroux and Gredt, 1972, MRID 05006363; Ambro-Balint, 1974, MRID 05013368; Snel and Edgington, 1970, MRID 05003663; Chin et al., 1970, MRID 05002177). Small amounts of carboxin sulfone (oxycarboxin) have been found in treated barley and wheat (Chin et al., 1970, MRID 05002177), and the p-hydroxylated derivative of carboxin has been identified in barley (Briggs et al., 1974, MRID 05002886). As crops mature, insoluble anilide complexes (these complexes of carboxin or carboxin derivatives with macromolecules such as lignin are insoluble in water and organic solvent and liberate aniline upon hydrolysis) increase (Briggs et al., 1974, 05002886; Snel and Edgington, 1970, MRID 05003663; Chin et al., 1970, MRID 05002177; Chin et al., 1972, MRID 00002941). Seven weeks after planting, acetone-insoluble residues were 23 percent of the total residue in barley and 40 percent in wheat (Chin et al., 1970, MRID 05002177). Polar metabolites of carboxin also increase during crop maturation (Leroux and

Gredt, 1972, MRID 05006363; Snel and Edgington 1970, MRID 05003663), but do not contribute significantly to the total residue in aerial portions of plants (Leroux and Gredt, 1972, MRID 05006363).

The uptake, distribution and metabolism of carboxin in plants has been adequately defined for the registered seed treatments. The residues of concern are: carboxin, carboxin sulfoxide, carboxin sulfone and insoluble anilide complexes.

2. Metabolism in Food-Producing Animals

Data on the metabolism of carboxin in food-producing animals have not been submitted to the Agency.

The Agency is not requiring an animal metabolism study to support the seed treatment use of carboxin. Data from a ruminant feeding study (see Residues in Animals) in conjunction with data on residues in crops (see Residues in Plants) indicate that total residues would be low or undetectable in tissues of animals fed commodities grown from seed treated with carboxin at application rates now permitted.

Additional metabolism and residue studies are being required to support the combined foliar and soil application to peanuts. If these studies show a higher residue level than that resulting from seed treatment, metabolism studies may be necessary on ruminants and poultry, since peanut meal, hulls, and soapstock are feed items.

3. Analytical Methods

A colorimetric method (Lane, 1970, MRID 05002737), by which carboxin and carboxin derivatives are determined as aniline, has been used to obtain most residue data on growing crops. Sensitivity of this method is 0.2 ppm.

Some residue data were obtained by an earlier version of the aforementioned colorimetric method (Lane, 1966, MRID 00003058).

A gas chromatographic method also based on determination of aniline has been used to gather data on residues in mature crops (Sisken and Newell, 1971, MRID 00003335). Sensitivity of the method is 0.2 ppm.

Recovery data for both methods are acceptable. No pesticide with tolerances on commodities on which carboxin is used has been found to interfere with the GLC method. Only pesticides hydrolyzable to 3,4-dichloroaniline, such as diuron and linuron, have been found to interfere with the colorimetric method (Lane and Sisken, 1977, MRID 00025467; Uniroyal Chemical, 1970, MRID 00025468; Uniroyal Chemical, 1977, MRID 00025483; Uniroyal Chemical, 1975, MRID 00002919; Uniroyal Chemical, 1975, MRID 00003054; Uniroyal Chemical, 1974, MRID 00002905; and Uniroyal Chemical, 1972, MRID 00002940).

The colorimetric method determines most carboxin derivatives including insoluble anilide complexes, but samples analyzed by the GLC method are not

likely to detect insoluble complexes. There have been no data submitted indicating the fraction of total residues determined by either method.

The colorimetric and GLC methods are subject to considerable interferences in untreated crop samples. A modified colorimetric method for residues hydrolyzable to aniline in meat, milk and eggs has been submitted (Uniroyal Chemical, 1973, MRID 00002857). The method differs from that of Lane in that samples undergo extraction and column-chromatography clean up steps prior to steam distillation. Sensitivity of the method is 0.2 ppm for meat and meat by-products, 0.05 ppm for eggs, and 0.02 - 0.05 ppm for milk. Untreated sample blanks are low, recovery data are adequate and other pesticide and pesticides with tolerances on meat, milk, and eggs do not interfere with the determination of carboxin. The method has been successfully tried out in an EPA laboratory and is published in the Pesticide Analytical Manual (PAM). This method has not been validated for commodities other than meat, milk and eggs.

In summary, both the colorimetric method of Lane and the GLC method of Sisken and Newell are suitable for obtaining residue data, although the latter method may not determine insoluble aniline complexes. Neither method is suitable for tolerance enforcement, but a modification of the colorimetric method is acceptable for enforcement of tolerances on meat, milk, and eggs. The modified method may be acceptable for tolerance enforcement on crops, if data, including data showing that the method determines insoluble anilide complexes, are submitted and the method was validated.

4. Residues in Plants

The following tables of residue data reflect the use of carboxin as a seed treatment. Residues in or on immature crops were analyzed primarily by the colorimetric method of Lane (1970, MRID 05002737), and residues in mature crops by the GLC method of Sisken and Newell (1971, MRID 00003335).

The residue level on carboxin-fortified plant samples did not change after seven months of subzero storage (colorimetric method, Stone, 1969, GS001201) or on fortified seed samples stored for 11 months at room temperature (GLC method, Sisken and Lane, 1970, MRID 00025466).

Corn: Residue data have been obtained on fodder (whole immature plants), forage (stalks at harvest), ears (without husks), and grain. Data were gathered in Illinois, Iowa, Indiana, Minnesota, Washington, Nebraska and North Carolina:

RESIDUE SUMMARY

SAMPLE	PHI (weeks)	RATE (s) (oz/cwt seed)	RESIDUE DISTRIBUTION ppm (as carboxin)		
			<0.2	0.2-0.5	>0.5
<u>DUST</u> 1/					
Fodder	4-21	1.5,3.0	12	0	0
Forage	22-23	1.0-3.0	6	0	0
Ears	21-25	1.0-3.0	8	0	0
grain	21-24	1.5-6.0	8	0	0
<u>READY-TO-USE (FLOWABLE)</u> 2/					
Fodder	4	1.4-2.7	0	1	1*
Fodder	5-9	1.4-2.7	11	1	0
Forage	15-26	0.7,4.0	10	0	0
Ears	15-26	0.7,4.0	10	0	0
<u>SOLUBLE CONCENTRATE-LIQUID</u> 3/					
Fodder	10-15	1.1	8	0	0
Forage	17-29	1.1	5	0	0
Ears	9-17	1.1	6	0	0
Grain	14-17	1.1	2	0	0

[*] (1.1 ppm)

^{1/} Uniroyal Chemical, 1977, MRID 00025483

^{2/} Uniroyal Chemical, 1973, MRID 00005852

^{3/} Uniroyal Chemical, 1975, MRID 00003356

COTTONSEED: Data on residues in whole cotton plants, seed, and in proceeded fractions of seed were obtained in California, Georgia, Mississippi, and Florida:

SAMPLE	PHI (weeks)	RATE (S) (oz/cwt seed)	RESIDUE DISTRIBUTION ppm (as carboxin)		
			<0.2	0.2-0.5	>0.5
<u>FLOWABLE FUNGICIDE</u> ^{1/}					
Seed	28-30	2.8-12.8	9	0	0
<u>WETTABLE POWDER</u> ^{2/,3/}					
Whole plant	2-4	3-12	0	0	13*
Whole plant	5-8	3-12	4	3	7**
Whole plant	10-14	3-12	6	2	0

[*] (2-32 ppm)

[**] (0.6-1.2 ppm)

^{1/} Uniroyal Chemical, 1973, MRID 00003129

^{2/} Uniroyal Chemical, 1970, MRID 00025468

^{3/} Uniroyal Chemical, 1967, MRID 00003185

Residues declined with a half-life of 5 days. Residues were less than 0.2 ppm in oil and meal from processed cottonseed grown from seed treated with 6 or 12 oz. carboxin per 100 lb. of seed (Sisken, 1970, MRID 00002938).

PEANUTS: Data on whole plants, peanut seed, meal and oil from processed seed, and on hulls were gathered from Georgia, North Carolina, Florida, and Texas:

SAMPLE	PHI (weeks)	RATE(s) (oz/cwt seed)	RESIDUE DISTRIBUTION ppm (as carboxin)		
			<0.2	0.2-0.5	>0.5
<u>WETTABLE POWDER</u> ^{1/} , ^{2/}					
Whole plant	2-4	4.5,9.0	0	2	8*
Whole plant	4-7	4.5,9.0	3	2	3**
Whole plant	8-10	4.5,9.0	4	2	2***

Residues in whole plants declined with a half-life of 4-5 days.

Hay ^{3/}	20	4.5,9.0	4	0	0
Hulls ^{4/}	20	4.5,9.0	4	0	0
Hulls ^{5/}	17	2.25,4.5	4****	0	0
Whole seed ^{5/}	17	2.25,4.5	4****	0	0
Meal ^{5/}	17	2.25,4.5	4****	0	0
Oil ^{5/}	17	2.25,4.5	4****	0	0

[*] (1.3-53 ppm)

[**] (1.3-3.6 ppm)

[***] (0.7-0.8 ppm)

[****] The formulation contained ¹⁴C-carboxin. All residues were less than 0.05 ppm.

^{1/} Uniroyal Chemical, 1970, MRID 00025468

^{2/} Uniroyal Chemical, 1969, MRID 00003045

^{3/} Uniroyal Chemical, 1974, MRID 00002905

^{4/} Uniroyal Chemical, 1974, MRID 00002903

^{5/} Collier et al., 1974, MRID 00003300

Additional residue studies are required to support the combined foliar and soil application to peanuts.

SMALL GRAINS: Data have been obtained in Idaho, Illinois, Utah, Kansas, North Dakota, Texas, and Manitoba on residues in whole plants and grain of oats, barley and wheat.

SAMPLE	PHI (weeks)	RATE(s) (oz/cwt seed)	RESIDUE DISTRIBUTION ppm (as carboxin)		
			<0.2	0.2-0.5	>0.5
<u>DUST</u>					
Wheat-whole plants ^{1/}	6	1.5,3.0	2	0	0
Wheat-grain ^{2/}	17-19	1.5,3.0	4	0	0
Wheat-straw ^{2/}	17-19	1.5,3.0	4	0	0
Barley-grain ^{3/}	18	1.5,3.0	2	0	0
Barley-straw ^{3/}	18	1.5,3.0	2	0	0
Oats-grain ^{4/}	23	1.5,3.0	2	0	0
Oats-straw ^{4/}	23	1.5,3.0	2	0	0
<u>READY-TO-USE (FLOWABLE)-^{5/}</u>					
Wheat-whole plants	4	1.2-4.0	2	2	6*
Wheat-whole plants	6-12	1.2-2.0	6	3	0
Wheat-whole plants	6-12	2.4-4.0	4	1	4**
Wheat-grain	12-40	1.2-5.6	12	0	0
Wheat-straw	12-40	1.2-4.0	10	0	0
Barley-whole plants	4	1.2-4.0	2	4	0
Barley-whole plants	6-12	1.2-2.4	7	1	0
Barley-whole plants	6-12	4.0	2	1	1#
Barley-grain	12-34	1.2-4.0	6	0	0
Barley-straw	12-34	1.2-4.0	6	0	0
<u>WETTABLE POWDER/DUST^{6/}</u>					
Wheat-whole plants ^{6/}	4-11	4	8	6	0
Wheat-whole plants ^{7/}	4-11	8	8	4	2##
Wheat-straw ^{7/}	14-39	3-6	8	0	0
Barley-whole plants ^{6/8/}	2-3	3-8	0	0	4\$
Barley-whole plants ^{8/}	4-11	3-4	14	4	0
Barley-whole plants ^{8/}	4-11	8	6	4	1##
Barley-grain ^{8/}	14-41	4-12	8	0	0
Oats-whole plants ^{8/}	6-11	4-12	9	3	0

Residues in whole plants declined with an approximate half-life of 10 days^{2,6,8/}

[*] (0.8-1.8 ppm)

[**] (0.6-0.7 ppm)

[#] (0.8 ppm)

[##] (0.6 ppm)

[\$] (0.9-13.3 ppm)

^{1/} Uniroyal Chemical, 1978, MRID 00003219

^{2/} Uniroyal Chemical, 1978, MRID 00003218

^{3/} Uniroyal Chemical, 1977, MRID 00003221

^{4/} Uniroyal Chemical, 1978, MRID 00003220

^{5/} Uniroyal Chemical, 1973, MRID 00003158

^{6/} Uniroyal Chemical, 1969, MRID 00003045

^{7/} Uniroyal Chemical, 1972, MRID 00002961

^{8/} Uniroyal Chemical, 1970, MRID 00025468

SORGHUM: Data were obtained on growing plants, grain, and fodder (plants after harvest) in Nebraska, South Dakota, Kansas, Missouri, and Texas.

SAMPLE	PHI (weeks)	RATE(s) (oz/cwt seed)	RESIDUE DISTRIBUTION ppm (as carboxin)		
			<0.2	0.2-0.5	>0.5
<u>READY-TO USE (FLOWABLE)</u> ^{1/}					
Whole plants	4-9	3,6	6	0	0
<u>SOLUBLE CONCENTRATE-LIQUID</u> ^{2/}					
Whole plants	4-14	1.1,2.2	22	2	0
Fodder	14	1.1,2.2	8	0	0
Grain	7-14	1.1,2.2	6	0	0
<u>WETTABLE POWDER/DUST</u> ^{1,3/}					
Whole plants	2-4	3,6	2	0	14*
Whole plants	4-11	3	8	6	0
Whole plants	4-11	6	7	1	6**

Residues declined with a half-life of 11 days ^{3/}

[*] (0.6-10.8)

[**] (0.6-1.7)

[1] Uniroyal Chemical, 1969, MRID 00003045

[2] Uniroyal Chemical, 1975, MRID 00003054

[3] Uniroyal Chemical, 1970, MRID 00025468

Data on residues in plants are adequate for the currently registered uses of carboxin, and are consistent with existing tolerances.

5. Residues of Carboxin in Animals

Three lactating cows were administered phenyl-¹⁴C-carboxin in the diet. Milk, urine, and feces were monitored daily for radioactivity, and after tendays of treatment the animals were sacrificed, and muscle, kidney, liver, and fat analyzed for radioactivity. Most of the radiolabel was excreted in the urine, the rate of excretion plateauing after the second day. Maximal radioactivity in the feces was reached in the first week. Trace radioactivity in milk plateaued after the first few days. Less than two percent of the ingested carboxin was detected in tissues at sacrifice. Residues were distributed as follows (Kennedy and Jenkins, 1971, MRID 00002945).

PPM ADMINISTERED	TOTAL RESIDUE (PPB - EXPRESSED AS CARBOXIN)				
	LIVER	KIDNEY	MUSCLE	FAT	MILK (MAXIMUM)
0.5	22	18	4	3	1
1.5	78	71	23	7	4
5.0	147	81	39	13	8

The feeding study adequately depicts the distribution of tagged residues in animals ingesting carboxin. However, the nature of the residue in food-producing animals is not known.

Data on residues in poultry and eggs have not been submitted to the Agency, however, the Agency is not requiring a poultry feeding study to support the seed treatment use of carboxin. The feeding study described above indicates no propensity for residues to accumulate in animal tissues, and data on residues in crops (see Residues in Plants) demonstrate the absence of detectable residues in poultry feed items.

B. LABELING REQUIREMENTS

Labels on all carboxin formulations are required to prohibit use of treated seed for food, feed, or oil purposes. A restriction on grazing livestock on treated areas for six weeks after planting should remain on products used on barley, corn, oats and wheat. Formulations used on cottonseed are required to bear a prohibition against grazing of livestock on treated areas and feeding hay grown from treated seed. The wettable powder formulation used on peanuts should continue to bear a restriction against grazing treated areas, but the current label restriction against feeding peanut hay (mature vines) to livestock is unnecessary because the established tolerance on peanut hay is adequate to cover seed treatment use.

C. SUMMARY OF DATA GAPS

Data demonstrating that the enforcement method is capable of determining the residue of concern in crops are required. Due to the SLN registrations for combined foliar and soil use, additional residue and metabolism studies on peanuts are required. If these studies show a higher residue level than that from seed treatment, residue and metabolism studies may be necessary on ruminants and poultry, since peanut meal, hulls, and soapstock are feed items.

VIII. ECOLOGICAL EFFECTS

- A. Ecological Effects Profile
- B. Ecological Effects Hazard Assessment
- C. Summary of Data Gaps

A. ECOLOGICAL EFFECTS PROFILE

Scientifically sound data on the toxicity of technical or end-use carboxin to nontarget organisms are very limited. One study (Fink, 1974, MRID 00003139) showed that technical carboxin is practically nontoxic to mallard ducks, with an LC_{50} value greater than 4640 ppm.

A few studies were presented which demonstrate the phytotoxicity of technical carboxin. A study showed that five ounces of technical carboxin per one hundred pounds of seed (cwt) as a seed treatment had no effect on wheat or barley (Kratka, 1971, MRID 05003530). Another study demonstrated that cotton seedlings grown in a nutrient solution containing 40 ppm carboxin suffered severe growth reduction (Davis, 1971, MRID 05001204). In a third study, a 100 ppm foliar drench of carboxin had no effect on begonia growth (Harris, 1975, MRID 05003951).

The following EC_{10} values for seed treatments were demonstrated using a 34% flowable concentrate formulation:

<u>Crop</u>	<u>Treatment</u>	<u>Reference</u>
Cotton	8 oz./cwt	Uniroyal, 1972, MRID 00003125
Wheat	28 oz./cwt	Uniroyal, 1973, MRID 00003094
Barley	28 oz./cwt	Uniroyal, 1973, MRID 00003094
Peanuts	2.2 lb a.i./acre*	Uniroyal, 1977, MRID 00003236

* This is a foliar spray application.

There was no effect on wheat or oats from a seed treatment of 12 oz. a.i./cwt, the highest level tested for the 25% dust formulation. The EC_{10} for barley was 3-4 oz. a.i./cwt for the 25% dust (Uniroyal, 1977, MRID 00005548).

There was no effect from using the 75% wettable powder formulations as seed treatments on the following crops:

<u>Crop</u>	<u>Treatment</u>	<u>Reference</u>
Peanuts	6 oz. a.i./cwt	Uniroyal, 1973, MRID 00002910
Corn	3 oz. a.i./cwt	Uniroyal, 1971, MRID 00002951
Wheat	8 oz. a.i./cwt	Clark, 1971, MRID 05003675
Barley	4 oz. a.i./cwt	Clark, 1971, MRID 05003675

Foliar applications of the wettable powder formulation at the rate of eight to nine pounds a.i./acre had only a slight effect on peanuts (Uniroyal, 1971, MRID 00002859 and Uniroyal, 1975, MRID 00003149).

There was no effect to peanuts resulting from the application of three pounds a.i./acre of 4% granular carboxin when applied at the pegging stage (Uniroyal, 1973, MRID 00002861).

There was no effect to cotton following a seed treatment using an experimental 10% liquid formulation at a rate of 8 oz./cwt (Uniroyal, 1974, MRID 00003001).

B. ECOLOGICAL EFFECTS HAZARD ASSESSMENT

Insufficient information is available on the technical, or end-use products, or their degradates to evaluate the potential impacts from carboxin to nontarget fish and wildlife species.

Due to the scarcity of information, a detailed plant hazard assessment cannot be made at this time. However, there is some information concerning the effects of carboxin on crops on which it is used. Those crops (corn, cotton, peanuts, wheat, barley, and oats) suffer no ill effects from seed treatments at the recommended rates, although cotton displayed phytotoxic symptoms when subjected to a root drench at a higher rate. Begonia, the only ornamental tested, suffered no injury from a foliar drench.

All existing uses could expose nontarget fish and wildlife species to the parent compound or its degradates, carboxin sulfoxide and carboxin sulfone. However, in the absence of appropriate data the significance of the exposure cannot be addressed.

C. SUMMARY OF DATA GAPS

The following tests are needed to meet minimum fish and wildlife data requirements to support the registration of carboxin:

Avian single-dose oral LD₅₀ test for either a wild waterfowl (preferably the mallard) or an upland game bird (preferably the bobwhite quail, other native

quail or the ring-necked pheasant). The species selected is to be the same as one of the two species selected for the avian dietary LC_{50} tests.

Avian dietary LC_{50} test for an upland game bird (preferably the bobwhite quail, other native quail, or the ringed-neck pheasant).

Fish acute LC_{50} test for one coldwater fish species, preferably the rainbow trout, and one warmwater species, preferably the bluegill.

Acute LC_{50} test for one species of freshwater invertebrate.

For the combined foliar and soil applications of carboxin, tests will need to be performed for the effect on seed germination, vegetative vigor, effects on algae, and aquatic macrophytes. These tests are not required for products registered solely for seed treatment.

Further data may be required depending upon the examination of the above data and adequate environmental chemistry data on the parent compound as well as its degradates.

IX. CASE BIBLIOGRAPHY

Guide to Use of This Bibliography

1. Content of Bibliography. This bibliography contains citations of all the studies reviewed by EPA in arriving at the positions and conclusions stated elsewhere in this standard. The bibliography is divided into two sections: (1) citations in numerical order that contributed information useful to the review of the chemical and are considered to be part of the data base supporting registrations under the standard, and (2) an alphabetical listing of all documents reviewed by the Agency. Primary sources for studies in this bibliography have been the body of data submitted to EPA and its predecessor agencies in support of past regulatory decisions, and the published technical literature.
2. Units of Entry. The unit of entry in this bibliography is called a "study". In the case of published materials, this corresponds closely to an article. In the case of unpublished materials submitted to the Agency, the Agency has sought to identify documents at a level parallel to a published article from within the typically larger volumes in which they were submitted. The resulting "studies" generally have a distinct title (or at least a single subject), can stand alone for purposes of review, and can be described with a conventional bibliographic citation. The Agency has attempted also to unite basic documents and commentaries upon them, treating them as a single study.
3. Identification of Entries. The entries in this bibliography are sorted by author, date of the document, and title. Each entry bears, to the left of the citation proper, a nine-digit numeric identifier. This number is unique to the citations and should be used at any time specific reference is required. This number is called the "Master Record Identifier" or "MRID". It is not related to the six-digit "Accession Number", which has been used to identify volumes of submitted data; see paragraph 4(3)(4) below for a further explanation. In a few cases, entries added to the bibliography late in the review may be preceded by a nine-character temporary identifier. This is also to be used whenever a specific reference is needed.
4. Form of the Entry. In addition to the Master Record Identifier (MRID), each entry consists of a bibliographic citation containing standard elements followed, in the case of materials submitted to EPA, by a description of the earliest known submission. The bibliographic conventions used reflect the standards of the American National Standards Institute (ANSI), expanded to provide for certain special needs. Some explanatory notes of specific elements follow:
 - a. Author. Whenever the Agency could confidently identify one, the Agency has chosen to show a personal author. When no individual was identified, the Agency has shown an identifiable laboratory or testing facility as author. As a last resort, the Agency has shown the first known submitter as author.

- b. Document Date. When the date appears as four digits with no question marks, the Agency took it directly from the document. When a four-digit date is followed by a question mark, the bibliographer deduced the date from evidence in the document. When the date appears as (19??), the Agency was unable to determine or estimate the date of the document.
- c. Title. This is the third element in the citation. In some cases it has been necessary for Agency bibliographers to create or enhance a document title. Any such editorial insertions are contained between square brackets.
- d. Trailing Parentheses. For studies submitted to us in the past, the trailing parentheses include (in addition to any self-explanatory text) the following elements describing the earliest known submissions:
 - (1) Submission Date. Immediately following the word 'received' appears the date of the earliest known submission, at the time that particular document was processed into the Pesticide Document Management System.
 - (2) Administrative Number. The next element, immediately following the word 'under', is the registration number, experimental permit number, petition number, or other administrative number associated with the earliest known submission, at the time that particular document was processed into the Pesticide Document Management System.
 - (3) Submitter. The third element is the submitter, following the phrase 'submitted by'. When authorship is defaulted to the submitter, this element is omitted.
 - (4) Volume Identification. The final element in the trailing parenthesis identifies the EPA accession number of the volume in which the original submission of the study appears. The six-digit accession number follows the symbol 'CDL', standing for "Company Data Library". This accession number is in turn followed by an alphabetic suffix which shows the relative position of the study within the volume. For example, within accession number 123456, the first study would be 123456-A; the second, 123456-B; the 26th, 123456-Z; and the 27th, 123456-AA.

OFFICE OF PESTICIDE PROGRAMS
REGISTRATION STANDARD NUMERICAL BIBLIOGRAPHY
Citations Considered to be Part of the Data Base Supporting
Registrations Under the Standard

MRID	CITATION
00002857	Uniroyal Chemical (1973) Enforcement Method for Vitavax® in Animal Tissue, Milk and Eggs. Method dated Sep 17, 1973. (Unpublished study received on unknown date under 3F1318; CDL:093547-D)
00002859	Uniroyal Chemical (1971) Reasoning in Support of the Petition: received Sep 27, 1972 under 3F1318; CDL:093547-O)
00002861	Uniroyal Chemical (1973) Vitavax® 10G: Peanuts: Phytotoxicity. (Unpublished study received Sep 28, 1976 under 400-130; CDL: 230405-N)
00002903	Uniroyal Chemical (1974) Residue Analysis of Peanut Hulls for Vitavax®. (Unpublished study received on unknown date under 4F1499; CDL:093979-C)
00002905	Uniroyal Chemical (1974) The Effects of EPA Tolerance Pesticides upon the Recovery of Vitavax® Residues from Peanuts. (Unpublished study received on unknown date under 4F1499; prepared in cooperation with Morse Laboratories, Inc.; CDL:093979-F)
00002910	Uniroyal Chemical (1973) Field Performance and Phytotoxicity Report. (Unpublished study received May 1, 1974 under 4F1499; CDL:094551-P)
00002935	Chin, W.T.; Dannals, L.E.; Kucharczyk, N. (1972) Environmental Fate Studies on Vitavax®: Status Report on PR 70-15. (Unpublished study received Mar 6, 1972 under 2F1191; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:093515-A)
00002938	Sisken, H.R. (1970) Determination of Residual Vitavax® in Meal and Oil from Cotton and Peanut Seed. (Unpublished study received Jul 28, 1972 under 0F0939; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091603-A)
00002941	Chin, W.T.; Kucharczyk, N.; Smith, A.E. (1972) Nature of Vitavax®-Derived Bound Residues in Plants. (Unpublished study received Oct 9, 1973 under 3F1318; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:092254-F)
00002943	Kennedy, G. (1971) Report to Uniroyal Chemical, Division of Uniroyal, Inc.: Distribution and Excretion Study with Vitavax® in Albino Rats: IBT No. E9289. (Unpublished study received Oct 9, 1973 under 3F1318; prepared by Industrial Bio-Test Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL: 092254-H)
00002944	Kennedy, G. (1971) Report to Uniroyal Chemical, Division of Uniroyal, Inc.: Characterization of 14C in Excreta of Rats Following Oral Administration of Vitavax®: IBT No. E9290. (Unpublished study received Oct 9, 1973 under 3F1318; prepared by Industrial Bio-Test Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:092254-I)

MRID	CITATION
00002945	Kennedy, G.; Jenkins, D.H. (1971) Report to Uniroyal Chemical, Division of Uniroyal, Inc.: Milk and Meat Residue Study in Dairy Cows Treated with 14C-Labeled Vitavax®: IBT No. J294. (Unpublished study received Oct 9, 1973 under 3F1318; prepared by Industrial Bio-Test Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:092254-J)
00002951	Uniroyal Chemical (1972) Field Performance Report. (Unpublished study including report, received Dec 23, 1972 under 3F1318; CDL:092253-F)
00002961	Uniroyal Chemical (1972) Residue Data: Vitavax®. (Unpublished study received on unknown date under 2F1191; prepared in cooperation with Morse Laboratories and Harris Laboratories; CDL:091003-A)
00002978	Uniroyal Chemical (1966) Vitavax® Assay Method. Method dated Jan 24, 1966. (Unpublished study received Aug 11, 1971 under 2F1191; CDL:093516-F)
00002995	Uniroyal Chemical (19??) Control Test AC-1045-D: Assay of Vitavax (and Its Formulations). Undated method. (Unpublished study received Jul 15, 1976 under 400-118; CDL:224929-C)
00003001	Uniroyal Chemical (1974) Phytotoxicity. (Unpublished study received Jul 15, 1976 under 400-118; CDL:224933-B)
00003030	Holsing, G.C. (1969) Final Report: Two-Year Dietary Administration--Dogs: Project No. 798-103. (Unpublished study received Jun 14, 1969 under 9G0819; prepared by Hazleton Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091420-E)
00003031	Holsing, G.C. (1969) Final Report: 24-Month Dietary Administration--Albino Rats: Project No. 798-102. (Unpublished study received Jun 14, 1969 under 9G0819; prepared by Hazleton Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091420-F)
00003032	Holsing, G.C. (1968) Final Report: Three-Generation Reproduction Study--Rats: Project No. 798-104. (Unpublished study received Jun 14, 1969 under 9G0819; prepared by Hazleton Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091420-G)
00003034	Binns, R.; Berczy, Z.; Clark, G.C. (1969) Acute Inhalation Toxicity to the Rat of Vitavax® Dust: 2540/68/417. (Unpublished study received Jun 14, 1969 under 9G0819; prepared by Huntingdon Research Centre, [Eng.] for Uniroyal, Ltd., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091420-K)

MRID	CITATION
00003035	Holsing, G.C. (1968) Final Report: Acute Eye Irritation--Rabbits: Project No. 798-151. (Unpublished study received Jun 14, 1969 under 9G0819; prepared by Hazleton Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091420-L)
00003044	Chin, W.T.; Stone, G.M.; Smith, A.E. (1969) 14C-Vitavax® Studies on Barley, Wheat and Cotton Plants. (Unpublished study received Jun 14, 1969 under 9G0819; Uniroyal Chemical, Bethany, Conn.; CDL:091420-W)
00003045	Uniroyal Chemical (1969) Disappearance Studies: Wheat and Barley; 1969 under 9G0819; CDL:091420-X)
00003054	Uniroyal Chemical (1975) The Effects of EPA Tolerance Pesticides upon the Recovery of Vitavax® Residues from Sorghum: (1) Grain, (2) Fodder and Forage. (Unpublished study including summary, received Jun 9, 1975 under 5F1638; prepared in cooperation with Morse Laboratories, Inc.; CDL:094948-G)
00003058	Lane, J.R. (1966) Residue Analysis for D-735 and/or F-461. Method dated May 16, 1966. (Unpublished study received Jan 25, 1967 under 400-81; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:121531-A)
00003063	Ozer, B.L. (1966) Report: Subacute (90-day) Feeding Studies with D-735 in Rats. (Unpublished study received Dec 14, 1966 under 400-EX-27; prepared by Food and Drug Research Laboratories, Inc. for United States Rubber Co., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:123429-F)
00003065	Carson, S. (1965) Approximate Acute Oral LD ₅₀ in Rats: Vitavax®. (Unpublished study received Dec 14, 1966 under 400-EX-27; prepared by Food and Drug Research Laboratories, Inc., for Naugatuck Chemical, Div. of United States Rubber Co., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:123429-N)
00003066	Carson, S. (1965) Approximate Acute Dermal LD ₅₀ in Rabbits: EX-27; prepared by Food and Drug Research Laboratories, Inc. for Naugatuck Chemical, Div. of United States Rubber Co., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:123429-O)
00003080	Babish, J.G. (1977) Acute Dermal Toxicity Study in Rabbits. (Unpublished study received Dec 29, 1977 under 400-107; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:232545-D)
00003081	Babish, J.G. (1977) Acute Oral Toxicity in Rats. (Unpublished study received Dec 29, 1977 under 400-107; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:232545-E)

MRID	CITATION
00003082	Babish, J.G. (1977) Acute Oral Toxicity in Rats. (Unpublished study received Dec 29, 1977 under 400-107; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:232545-F)
00003083	Babish, J.G. (1977) Acute Inhalation Study in Rats. (Unpublished study received Dec 29, 1977 under 400-107; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:232545-G)
00003084	Uniroyal Chemical (1976) Vitavax® Manufacturing Process. (Unpublished study including letter dated Sep 9, 1976 from T.E. Geise to John Pryzbylek, received Sep 13, 1976 under 400-129; CDL: 225601-A)
00003087	Cardona, R.A.; Dannals, L.E.; Smilo, A.R. (1976?) Fate of Vitavax (R)? in Soil: Project No. 7511. (Unpublished study received May 3, 1977 under 400-80; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:231932-B)
00003088	Smilo, A.R.; Lacadie, J.A.; Cardona, R. (1977) Photochemical Fate of Vitavax® in Solution. (Unpublished study received May 3, 1977 under 400-80; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:231932-C)
00003094	Uniroyal Chemical (1973) Field Performance Report: UNI-1088 and 1090. (Unpublished study received Jun 26, 1973 under 400-108; CDL:008327-F)
00003114	Dannals, L.E.; Campbell, C.R.; Cardona, R.A. (1976) Environmental Fate Studies on Vitavax®: Status Report II on PR 70-15. Includes three undated methods. (Unpublished study received Mar 17, 1976 under 400-80; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:223866-A)
00003116	Babish, J.G. (1977) Acute Inhalation Study in Rats. (Unpublished study received Jan 24, 1978 under 400-81; prepared by Food and Drug Research Laboratories, Inc.; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:233500-B)
00003118	Brusick, D.J. (1977) Mutagenicity Evaluation of D735: Final Report: LBI Project No. 2683. (Unpublished study received Jan 24, 1978 under 400-81; prepared by Litton Bionetics, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:233500-D)
00003119	Babish, J.G. (1977) Primary Skin Irritation Study with Rabbits. (Unpublished study received Jan 24, 1978 under 400-81; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:233500-E)

MRID	CITATION
00003120	Knickerbocker, M. (1977) Teratologic Evaluation of Vitavax® Technical in Sprague Dawley Rats. (Unpublished study received Jan 24, 1978 under 400-81; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:233500-F)
0003125	Uniroyal Chemical (1972) Field Performance Report. (Unpublished study received May 16, 1973 under 400-107; CDL:003284-H)
00003129	Uniroyal Chemical (1973) Residues in PPM. (Unpublished study received May 16, 1973 under 400-107; prepared in cooperation with Morse Laboratories, Inc.; CDL:003284-P)
00003139	Fink, R. (1974) Final Report: Eight-Day Dietary LC ₅₀ -Mallard Ducks: Project No. 117-102. (Unpublished study including official analytical report, received Mar 11, 1974 under 400-106; prepared by Truslow Farms, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:128723-A)
00003149	Uniroyal Chemical (1975) Field Evaluation Report. (Unpublished study received Sep 13, 1976 under 400-129; CDL:225604-W)
00003152	Holsing, G.C. (1969) Final Report: 24-Month Dietary Administration—Albino Rats: Project No. 798-102. (Unpublished study received Oct 21, 1971 under 400-90; prepared by Hazleton Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:050926-A)
00003158	Uniroyal Chemical (1973) Vitavax Flowable Fungicide: Summary of Residues in Wheat and Barley. (Unpublished study received Dec 5, 1973 under 400-107; CDL:023365-S)
00003161	Babish, J.G. (1976) Rabbit Eye Irritation Study. (Unpublished study received Apr 21, 1977 under unknown admin. no.; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:229370-C)
00003162	Babish, J.G. (1976) Primary Skin Irritation Study with Rabbits. (Unpublished study received Apr 21, 1977 under unknown admin. no.; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:229370-D)
00003172	Puchalsky, C.B. (1968) Infrared Method for the Assay of D-735, and Its Content in Formulations. Method dated Jul 15, 1968. (Unpublished study received Mar 25, 1969 under 9G0819; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:093520-H)
00003185	Uniroyal Chemical (1967) Residue Analysis of Cotton Seedlings from Vitavax®-Treated Cotton Seeds. (Unpublished study received Nov 29, 1967 under 400-EX-28; CDL:123430-A)

MRID	CITATION
00003216	Holsing, G.C. (1968) Final Report: Repeated Dermal Application-- Rabbits: Project No. 798-148. (Unpublished study received Jan 13, 1969 under 400-EX-33; prepared by Hazleton Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL: 123439-M)
00003218	Uniroyal Chemical (1978) Residues in PPM: Wheat: Vitavax® 25DB: Grain & Straw. (Unpublished study received May 10, 1978 under 400-115; prepared in cooperation with Morse Laboratories, Inc.; CDL:235653-B)
00003219	Uniroyal Chemical (1978) Residues in PPM: Wheat: Vitavax® 25DB: Forage. (Unpublished study received May 10, 1978 under 400-115; prepared in cooperation with Morse Laboratories, Inc.; CDL: 235653-C)
00003220	Uniroyal Chemical (1978) Residues in PPM: Oats: Vitavax® 25DB: Grain & Straw. (Unpublished study received May 10, 1978 under 400-115; prepared in cooperation with Morse Laboratories, Inc.; CDL:235653-D)
00003221	Uniroyal Chemical (1977) Residues in PPM: Barley: Vitavax® 25DB: Grain & Straw. (Unpublished study received May 10, 1978 under 400-115; prepared in cooperation with Morse Laboratories, Inc.; CDL:235653-E)
00003224	Uniroyal Chemical (1978) Response to EPA Letter Dated 6 October, 1977: Environmental Fate of Vitavax® Fungicide. Summary of studies 236662-F through 236662-J. Includes method no. 2344 dated Aug 22, 1978. (Unpublished study received Dec 27, 1978 under 400-80; prepared in cooperation with Morse Laboratories, Inc.; CDL:236662-A)
00003225	Dzialo, D.G.; Lacadie, J.A. (1978) Aerobic Soil Study of 14C-Vitavax® in Sandy Soil: Project No. 7746-1. (Unpublished study received Dec 27, 1978 under 400-80; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:236662-F)
00003226	Dzialo, D.G.; Lacadie, J.A.; Cardona, R.A. (1978) Anaerobic Soil Metabolism of 14C-Vitavax® in Sandy Soil: Project No. 7751. (Unpublished study received Dec 27, 1978 under 400-80; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:236662-G)
00003229	Lacadie, J.A.; Dzialo, D.G.; Dannals, L.E. (1978) Vitavax® 14C Aged Laboratory Column Leaching in Sandy Soil: Project No. 7832. (Unpublished study received Dec 27, 1978 under 400-80; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:236662-J)
00003231	Uniroyal Chemical (197?) Vitavax®-Flowable Fungicide: Vitavax®-3F Fungicide: Basic Manufacturing Process. (Unpublished study received Mar 20, 1979 under 400-80; CDL:098014-B)

MRID	CITATION
00003236	Uniroyal Chemical (1977) Vitavax®-3F: Peanuts. (Unpublished study received May 30, 1978 under 400-EX-55; CDL:234122-B)
00003296	Uniroyal Chemical (19??) Vitavax®: Chemical and Physical Properties of Vitavax® Technical. (Unpublished study received Sep 27, 1972 under 3F1318; CDL:093547-J)
00003300	Collier, Y.J.; Womer, J.M.; Stone, G.M.; Kucharczyk, N. (1974) Data of Vitavax®-14C Derived Residues in Peanut Hulls, Seed, Oil and Meal. (Unpublished study received on unknown date under 4F1499; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:093979-E)
00003301	Bailey, D.E. (1976) Rabbit Eye Irritation Study: Vitavax® 75W. (Unpublished study received Sep 13, 1976 under 400-129; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:225606-D)
00003311	Matthews, R.J. (1970) Primary Skin Irritation--Rabbits: Vitavax®: Final Report. (Unpublished study received May 8, 1972 under 2F1191; prepared by Pharmakon Laboratories, submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091003-G)
00003314	Matthews, R.J. (1970) Acute Dermal Application--Rabbits: Vitavax®: Final Report. (Unpublished study received May 8, 1972 under 2F1191; prepared by Pharmakon Laboratories, submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091003-J)
00003317	Matthews, R.J. (1970) Acute LD ₅₀ —Rats (Oral): Vitavax®: Final Report. (Unpublished study received May 8, 1972 under 2F1191; prepared by Pharmakon Laboratories, submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091003-M)
00003335	Sisken, H.R.; Newell, J.E. (1971) Determination of residues of Vitavax® and its sulfoxide in seeds. Journal of Agricultural and Food Chemistry 19(4):738-741. (Also In unpublished submission received Apr 21, 1977 under unknown admin. no.; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:229370-P)
00003356	Uniroyal Chemical (1975) Residues of Vitavax®--EVS Concentrate in Corn. (Unpublished study received Jun 10, 1975 under 400-124; prepared in cooperation with Morse Laboratories, Inc.; CDL:235928-C)
00005540	Spare, W. (1979) Report: Vitavax® Microbial Metabolism in Soil and Its Effect on Microbes. Includes undated method. (Unpublished study received Mar 28, 1979 under 400-81; prepared by Biospherics, Inc. in cooperation with United States Testing Co., Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:098029-A)

MRID	CITATION
00005544	Kuc, W.J.; Doebbler, G.F. (1979) Uniroyal Chemical 14C-Vitavax® Bluegill Sunfish Bioconcentration Study: UCES Project No. 11506-82. Includes undated method. (Unpublished study including metabolite study, received Mar 28, 1979 under 400-81; prepared by Union Carbide Corp., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:098029-F)
00005545	Kuc, W.J.; Doebbler, G.F. (1979) Uniroyal Chemical 14C-Vitavax® Bioconcentration Study: Channel Catfish, <u>Ictalurus punctatus</u> (Rafinesque): Project No. 11506-82. Includes undated method. (Unpublished study including metabolite study, received Mar 28, 1979 under 400-81; prepared by Union Carbide Corp., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:098029-G)
G 005548	Uniroyal Chemical (1977) Vitavax®-25DB: Phytotoxicity. (Unpublished study received Feb 28, 1978 under 400-115; prepared in cooperation with the University of Idaho, Experiment Station, CDL:238076-D)
00005852	Uniroyal Chemical (1973) Residues in PPM: Corn: UNI-1088. (Unpublished study received Mar 28, 1977 under 400-107; prepared in cooperation with New York State Univ.--Oswego, Lake Ontario Environmental Laboratory and Morse Laboratories, Inc.; CDL: 238083-A)
00005856	Stevens, K.R. (1979) Primary Skin Irritation Study with Rabbits. (Unpublished study received Feb 28, 1979 under 400-124; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:237477-B)
00005857	Stevens, K.R. (1979) Report: Eye Irritation Test in Rabbits. (Unpublished study received Feb 28, 1979 under 400-124; prepared by Food & Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:237477-C)
00005858	Stevens, K.R. (1979) Report: Acute Dermal Toxicity Study in Rabbits. (Unpublished study received Feb 28, 1979 under 400-124; prepared by Food & Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:237477-D)
00005859	Uniroyal Chemical (1977) Vitavax® Fungicide: Technical Data Sheet. (Unpublished study received Feb 6, 1979 under 400-EX-58; CDL:237329-A)
00025466	Sisken, H.R.; Lane, J.R. (1970) Storage Stability Studies of Vitavax® Residue in Grain. (Unpublished study received Jan 12, 1970 under 0F0939; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091604-E)
00025468	Lane, J.R.; Sisken, H.R. (1970) Specificity of the Vitavax® Method for Seeds. (Unpublished study received Jan 12, 1970 under 0F0939; submitted by Uniroyal Chemical, Bethany, Conn.; CDL: 091604-G)

MRID	CITATION
00025483	Uniroyal, Incorporated (1974) Summary of Corn and Rice Residue Data. (Unpublished study received Jun 17, 1974 under 5F1525; prepared in cooperation with Morse Laboratories, Inc. and State Univ. College of New York--Oswego, Lake Ontario Environmental Laboratory; CDL:094043-H)
05001172	Verma, R.K.; Vyas, S.C. (1976) Uptake, translocation and persistence of five systemic fungicides in gram seedlings. <i>Pesticides</i> 10(12):21-24.
05001302	Thapliyal, P.N.; Sinclair, J.B. (1970) Uptake of three systemic fungicides by germinating soybean seed. <i>Phytopathology</i> 60(9):1373-1375.
05001304	Thapliyal, P.N.; Sinclair, J.B. (1971) Translocation of benomyl, carboxin, and chloroneb in soybean seedlings. <i>Phytopathology</i> 61(10):1301-1302.
05002176	Chin, W.T.; Stone, G.M.; Smith, A.E. (1970) Degradation of carboxin (Vitavax®) in water and soil. <i>Journal of Agricultural and Food Chemistry</i> 18(4):731-732.
05002177	Chin, W.T.; Stone, G.M.; Smith, A.E. (1970) Metabolism of carboxin (Vitavax®) by barley and wheat plants. <i>Journal of Agricultural and Food Chemistry</i> 18(4):709-712.
05002737	Lane, J.R. (1970) Colorimetric microdetermination of Vitavax® (5,6 dihydro-2-methyl-1, 4-oxathiin-3-carboxanilide) residues in crops. <i>Journal of Agricultural and Food Chemistry</i> 18(3):409-412.
05002793	Bolkan, H.A.; Milne, K.S. (1975) Systemic uptake of four fungicides by potato plants. <i>Plant Disease Reporter</i> 59(3):214-218.
05002823	Buchenauer, H. (1975) Differences in light stability of some carboxylic acid anilide fungicides in relation to their applicability for seed and foliar treatment. <i>Pesticide Science</i> 6(5):525-535.
05002886	Briggs, D.E.; Waring, R.H.; Hackett, A.M. (1974) The metabolism of carboxin in growing barley. <i>Pesticide Science</i> 5(5):599-607.
05003218	El-Dib, M.A.; Aly, O.A. (1976) Persistence of some phenylamide pesticides in the aquatic environment--III. Biological degradation. <i>Water Research</i> 10(12):1055-1059.
05003663	Snel, M.; Edgington, L.V. (1970) Uptake, translocation and decomposition of systemic oxathiin fungicides in bean. <i>Phytopathology</i> 60(12):1708-1716.
05003664	Kirk, B.T.; Sinclair, J.B.; Lambremont, E.N. (1969) Translocation of 14C-labeled chloroneb and DMOC in cotton seedlings. <i>Phytopathology</i> (59):1473-1476.

MRID	CITATION
05003673	Berggren, G.T.; Pinckard, J.A. (1973) Movement of the systemic fungicide carboxin into cotton seedlings assisted by the systemic insecticide monocrotophos. <i>Plant Disease Reporter</i> 57(8):642-645.
05003675	Clark, R.V. (1971) Influence of Vitavax® seed treatment and loose smut infection on yields of barley and wheat. <i>Canadian Plant Disease Survey</i> 51(3):131-133.
05003778	Handa, S.K. (1978) Spectrophotometric determination of carboxin in formulations after extraction and thin layer chromatographic separation. <i>Journal of the Association of Official Analytical Chemists</i> 61(4):971-972.
05003852	Lyr, H.; Ritter, G.; Banasiak, L. (1974) Detoxication of carboxin. <i>Zeitschrift fuer Allgemeine Mikrobiologie</i> 14(4):313-320.
05003915	El-Dib, M.A.; Aly, O.A. (1976) Persistence of some phenylamide pesticides in the aquatic environment--I. Hydrolysis--II. Adsorption on clay minerals--III. Biological degradation. <i>Water Research</i> 10(12):1047-1053,1055-1059.
05003947	Curley, R.L.; Burton, J.C. (1975) Compatibility of <u>Rhizobium japonicum</u> with chemical seed protectants. <i>Agronomy Journal</i> 67(6):807-808.
05004129	Michail, S.H.; Elarosi, H.; Abd-El-Rehim, M.A.; Shohda, W.T. (1975) Persistence of certain systemic and non-systemic fungicides in the soil, with special reference to the method of application. <i>Phytopathologia Mediterranea</i> 14(2/3):138-139.
05004996	Chin, W.T.; Stone, G.M.; Smith, A.E.; Schmeling, B. von (1970) Fate of carboxin in soil, plants and animals. Pages 322-327, In <i>Proceedings of the Fifth British Insecticide and Fungicide Conference</i> ; Nov 17-20, 1969, Brighton, England. Vol. 2. Brighton, England: British Insecticide and Fungicide Conference.
05005076	Day, E.W., Jr.; Koons, J.R.; Ballee, D.L.; Duane, W.C.; Stallard, D.E.; Wolfe, A.L.; Lynch, V.P.; Edwards, M.J.; Wood, J.S., Jr.; Nishi, K.; Konishi, K.; Stone, G.M. (1976) Fungicides. Pages 251-261, 263-283, 285-297, 299-331, In <i>Analytical Methods for Pesticides and Plant Growth Regulators</i> . Vol. 8: Government Regulations, Pheromone Analysis, Additional Pesticides. New York: Academic Press.
05006363	Leroux, P.; Gredt, M. (1972) Devenir de la carboxine utilisee en enrobage de semences de ble: absorption, transport et degradation. [The fate of carboxin used as a dressing for wheat seed: absorption, translocation and degradation.] <i>Phytiatrie-Phytopharmacie</i> 21(1):45-60.

MRID	CITATION
05013368	Balintne Ambro, I. (1974) Tracerindikacios vizsgalatok a [6-14C] Vitavax fungicid metabolizmusanak tanulmanyozasara. [Uptake, translocation and decomposition of the fungicide Vitavax-[6-14C].] Magyar Kemikusok Lapja. [Journal of the Hungarian Chemists.] XXIX(8):393-401.
GS001201	Stone, G.M. (1969) Stability of Vitavax® in Stored Plants. (Unpublished study received Jan 12, 1970 under 0F0939; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:09604-A)
GS001202	Babish, J.G. (1976) Eye Irritation in rabbits- Vitavax®-17. (Unpublished study received June 20, 1978 under 400-113; prepared by Food and Drug Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:234185-A)
GS001203	Babish, J.G. (1977) Acute Dermal LD ₅₀ in rabbits - Vitavax®-17. (Unpublished study received June 20, 1978 under 400-113; prepared by Food and Drug Laboratories, Inc. submitted by Uniroyal Chemical, Bethany, Conn.; CDL:234185-B)
GS001204	Babish, J.G. (1977) Acute Inhalation LD ₅₀ in rats- Vitavax®-17. (Unpublished study received June 20, 1978 under 400-113; prepared by Food and Drug Laboratories, Inc. submitted by Uniroyal Chemical, Bethany, Conn.; CDL:234185-C)
GS001205	Babish, J.G. (1977) Acute Oral LD ₅₀ in rats- Vitavax®-17. (Unpublished study received June 20, 1978 under 400-113; prepared by Food and Drug Laboratories, Inc. submitted by Uniroyal Chemical, Bethany, Conn.; CDL:234185-D)
GS001206	Babish, J.G. (1977) Skin Irritation in rabbits - Vitavax®-17. (Unpublished study received June 20, 1978 under 400-113; prepared by Food and Drug Laboratories, Inc. submitted by Uniroyal Chemical, Bethany, Conn.; CDL:234185-E)

OFFICE OF PESTICIDE PROGRAMS
REGISTRATION STANDARD ALPHABETICAL BIBLIOGRAPHY
Listing of All Documents Reviewed by the Agency

MRID	CITATION
05006895	Agnihotri, V.P. (1971) Effects of certain fungitoxicants on the viability and pathogenicity of sclerotia of <u>Waitea circinata</u> . Phytopathologische Zeitschrift 70(1):71-80.
05004349	Allam, A.I.; Schilling, P.E.; Sinclair, J.B. (1969) Uptake of Vitavax (DMO) by germinating cottonseed. Phytopathology 59(2):265-266.
05004123	Allam, A.I.; Sinclair, J.B. (1968) Uptake and movement of Vitavax® by germinating cottonseed. Phytopathology 58(8):1041.
05002174	Allam, A.I.; Sinclair, J.B. (1969) Degradation of DMOC (Vitavax) in cotton seedlings. Phytopathology 59(10):1548-1549.
00003322	Allam, A.I.; Sinclair, J.B. (1969) Degradation of DMOC (Vitavax) in cotton seedlings. Phytopathology 59(10):1548-1549; Taken from Pestdoc 2(34):86741J. (Abstract no. 886/093; also In unpublished submission received Jan 12, 1970 under OF0939; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:093245-B)
05003381	Andreeva, E.I.; Usmanov, M.T.; Kurganova, L.V. (1972) Metodika ispytaniya preparatov protiv kornevoi gnili. [A method of testing preparations against root rot.] Khlopkovodstvo. [Cotton Growing.] (11):19-21.
00025473	Arny, D.C.; Blackmon, C.W.; Hansing, E.D.; et al. (1969) [Efficacy Data for Vitavax on Cereal and Forage Crops]. (Unpublished study received Jan 12, 1970 under OF0939; prepared by Univ. of Wisconsin and others, submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091604-L)
00003162	Babish, J.G. (1976) Primary Skin Irritation Study with Rabbits. (Unpublished study received Apr 21, 1977 under unknown admin. no.; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:229370-D)
00003161	Babish, J.G. (1976) Rabbit Eye Irritation Study. (Unpublished study received Apr 21, 1977 under unknown admin. no.; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:229370-C)
00003014	Babish, J.G. (1976) Vitavax®200 Flowable Fungicide: 076008: Primary Skin Irritation Study with Rabbits. (Unpublished study received Jul 14, 1978 under 400-112; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:235042-F)
00003013	Babish, J.G. (1976) Vitavax®200 Flowable Fungicide: 076008: Rabbit Eye Irritation Study. (Unpublished study received Jul 14, 1978 under 400-112; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL: 235042-E)

MRID	CITATION
GS001202	Babish, J.G. (1976) Eye irritation in rabbits-Vitavax®17. (Unpublished study received June 20, 1978 under 400-113; prepared by Food and Drug Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:234185-A)
00003080	Babish, J.G. (1977) Acute Dermal Toxicity Study in Rabbits. (Unpublished study received Dec 29, 1977 under 400-107; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:232545-D)
00021621	Babish, J.G. (1977) Acute Dermal Toxicity Study in Rabbits. (Unpublished study received Apr 25, 1979 under 400-136; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:238226-E)
GS001203	Babish, J.G. (1977) Acute dermal LD ₅₀ , rabbits-Vitavax®17. (Unpublished study received June 20, 1978 under 400-113; prepared by Food and Drug Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:234185-B)
00003083	Babish, J.G. (1977) Acute Inhalation Study in Rats. (Unpublished study received Dec 29, 1977 under 400-107; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:232545-G)
00003116	Babish, J.G. (1977) Acute Inhalation Study in Rats. (Unpublished study received Jan 24, 1978 under 400-81; prepared by Food and Drug Research Laboratories, Inc.; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:233500-B)
00021622	Babish, J.G. (1977) Acute Inhalation Study in Rats. (Unpublished study received Apr 25, 1979 under 400-136; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:238226-F)
00029650	Babish, J.G. (1977) Acute Inhalation Study in Rats: Laboratory No. 5483. (Unpublished study received Apr 25, 1979 under 400-136; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:238226-F)
GS001204	Babish, J.G. (1977) Acute Inhalation in rats-Vitavax®17. (Unpublished study received June 20, 1978 under 400-113; prepared by Food and Drug Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:234185-C)
00003082	Babish, J.G. (1977) Acute Oral Toxicity in Rats. (Unpublished study received Dec 29, 1977 under 400-107; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:232545-F)
00003081	Babish, J.G. (1977) Acute Oral Toxicity in Rats. (Unpublished study received Dec 29, 1977 under 400-107; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:232545-E)

MRID	CITATION
GS001205	Babish, J.G. (1977) Acute Oral LD ₅₀ in rats-Vitavax®17. (Unpublished study received June 20, 1978 under 400-113; prepared by Food and Drug Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:234185-D)
00021620	Babish, J.G. (1977) Approximate Acute Oral Toxicity (LD ₅₀) in Rats. (Unpublished study received Apr 25, 1979 under 400-136; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:238226-D)
00003119	Babish, J.G. (1977) Primary Skin Irritation Study with Rabbits. (Unpublished study received Jan 24, 1978 under 400-81; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:233500-E)
00021619	Babish, J.G. (1977) Primary Skin Irritation Study with Rabbits. (Unpublished study received Apr 25, 1979 under 400-136; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:238226-C)
00005866	Babish, J.G. (1977) Report: Acute Dermal Toxicity Study in Rabbits. (Unpublished study received Feb 6, 1979 under 400-EX-58; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:237329-I)
00005549	Babish, J.G. (1977) Report: Acute Oral Toxicity in Rats. (Unpublished study received Feb 6, 1979 under 400-EX-58; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:237329-H)
00005864	Babish, J.G. (1977) Report: Primary Skin Irritation Study with Rabbits. (Unpublished study received Feb 6, 1979 under 400-EX-58; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:237329-F)
GS001206	Babish, J.G. (1977) Skin irritation in rabbits-Vitavax®17. (Unpublished study received June 20, 1978 under 400-113; prepared by Food and Drug Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:234185-E)
00003015	Babish, J.G. (1977) Vitavax®200 Flowable Fungicide: G 037132 JS; BL 8538: Acute Dermal Toxicity Study in Rabbits. (Unpublished study received Jul 14, 1978 under 400-112; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:235042-G)
00003011	Babish, J.G. (1977) Vitavax®200 Flowable Fungicide: BL #8229: Acute Inhalation Study in Rats. (Unpublished study received Jul 14, 1978 under 400-112; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:235042-C)

MRID	CITATION
05020737	Bachofer, R. (1976) Mikrobieller Abbau von Sacureanilid-Fungiziden. [Microbial breakdown of acid anilide fungicides.] Zentrablatt fuer Bakteriologie, Parasitenkunde, Infektionskrankheiten und Hygiene. Abteilung 1 Originale. Reihe B 162(1/2):153-156.
05005110	Bachofer, R.; Oltmanns, O.; Lingens, F. (1973) Isolation and characterization of a <i>Nocardia</i> -like soil-bacterium, growing on carboxanilide fungicides. Archiv fuer Mikrobiologie 90(2):141-149.
05003943	Backman, P.A. (1977) Comparative fungicide sensitivity within the cowpea inoculation group of <i>Rhizobium</i> . Pages 221, In Proceedings of the American Phytopathological Society. Vol. 4. St. Paul, Minn. American Phytopathological Society.
00003163	Bailey, D.E. (1976) Rabbit Eye Irritation Study. (Unpublished study received Apr 21, 1977 under unknown admin. no.; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:229370-E)
00021618	Bailey, D.E. (1976) Rabbit Eye Irritation Study. (Unpublished study received Apr 25, 1979 under 400-136; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:238226-B)
00003302	Bailey, D.E. (1976) Rabbit Eye Irritation Study: Vitavax [®] HB-10. (Unpublished study received Sep 28, 1976 under 400-118; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:230403-B)
00003301	Bailey, D.E. (1976) Rabbit Eye Irritation Study: Vitavax [®] 75WP. (Unpublished study received Sep 13, 1976 under 400-129; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:225606-D)
00030103	Bailey, D.E. (1976) Rabbit Eye Irritation Study: Laboratory No. 5186c. (Unpublished study received Apr 25, 1979 under 400-136; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:238226-B)
00005865	Bailey, D.E. (1976) Report. Rabbit Eye Irritation Study. (Unpublished study received Feb 6, 1979 under 400-EX-58; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:237329-G)
00003005	Bailey, D.E. (1976) 10G Vitavax [®] : Rabbit Eye Irritation Study. (Unpublished study received Sep 28, 1976 under 400-130; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:230409-D)

MRID

CITATION

- 05020699 Bailey R.A. (1979) Possibilities for the control of sugarcane smut (Ustilago scitaminea) with fungicides. Pages 137-142, In Proceedings of the Annual Congress of the South African Sugar Technologists' Association. Vol. 53. Mount Edgecombe, Natal, South Africa: South African Sugar Technologists' Association.
- 05002489 Baker, P.B.; Farrow, J.E.; Hoodless, R.A. (1973) Agricultural fungicides: I. Identification of systemic fungicides using thin-layer chromatography. Journal of Chromatography 81(1):174-177.
- 05002749 Baker, P.B.; Hoodless, R.A. (1974) Analytical methods for the detection and determination of residues of systemic fungicides. Pesticide Science 5(4):465-472.
- 05006289 Balasubramanya, R.H.; Patil, R.B. (1976) Degradation of carboxin and oxycarboxin by a species of Pseudomonas isolated from soil. Madras Agricultural Journal 63(8/10):505-508.
- 05013368 Balintne Ambro, I. (1974) Tracerindikacios vizsgalatok a [6-14C] Vitavax fungicid metabolizmusanak tanulmanyozasara. [Uptake, translocation and decomposition of the fungicide Vitavax-[6-14C].] Magyar Kemikusok Lapja [Journal of the Hungarian Chemists.] XXIX(8):393-401.
- 00003337 Bell, D.K. (1967) Peanut Seed Treatment Tests 1967. (Unpublished study received Feb 23, 1968 under 400-EX-33; prepared by Georgia Coastal Plain Experiment Station, Dept. of Plant Pathology, submitted by Uniroyal Chemical, Bethany, Conn.; CDL:123438-C)
- 05003600 Bhaktavatsalam, G.; Tripathi, R.K. (1975) Uptake, translocation and degradation of carboxin and oxycarboxin in pearl millet seeds. Indian Journal of Mycology and Plant Pathology 5(1):110-111.
- 00003034 Binns, R.; Berczy, Z.; Clark, G.C. (1969) Acute Inhalation Toxicity to the Rat of Vitavax® Dust: 2540/68/417. (Unpublished study received Jun 14, 1969 under 9G0819; prepared by Huntingdon Research Centre, [Eng.] for Uniroyal, Ltd., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091420-K)
- 05009413 Bland, C.E.; Ruch, D.G.; Salser, B.R.; Lightner, D.V. (1976) Chemical Control of Lagenidium, a Fungal Pathogen of Marine Crustacea. Raleigh, N.C.: University of North Carolina Sea Grant Program. (Sea Grant publication no. UNC-SG-76-02; available from: NTIS, Springfield, VA; PB-257 346)
- 05002793 Bolkan, H.A.; Milne, K.S. (1975) Systemic uptake of four fungicides by potatoe plants. Plant Disease Reporter 59(3):214-218.
- 05012098 Born, G.L. (1974) Root infection of woody hosts with Verticillium albo-atrum. Illinois Natural History Survey Bulletin 31(6):209-248.

MRID	CITATION
05002886	Briggs, D.E.; Waring, R.H.; Hackett, A.M. (1974) The metabolism of carboxin in growing barley. <i>Pesticide Science</i> 5(5):599-607.
00003118	Brusick, D.J. (1977) Mutagenicity Evaluation of D735: Final Report: LBI Project No. 2683. (Unpublished study received Jan 24, 1978 under 400-81; prepared by Litton Bionetics, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:233500-D)
05002823	Buchenauer, H. (1975) Differences in light stability of some carboxylic acid anilide fungicides in relation to their applicability for seed and foliar treatment. <i>Pesticide Science</i> 6(5):525-535.
05012613	Buchman, O.; Pri-Bar, I.; Shimoni, M.; Smoskowitz, L. (1974) Application d'une methode non specifique de tritiation de molecules organiques. [Application of a nonspecific method of tritiation of organic molecules.] <i>Journal of Labelled Compounds (Brussels)</i> X(2):345-354.
00003087	Cardona, R.A.; Dannals, L.E.; Smilo, A.R. (1976?) Fate of Vitavax® in Soil: Project No. 7511. (Unpublished study received May 3, 1977 under 400-80; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:231932-B)
00003066	Carson, S. (1965) Approximate Acute Dermal LD ₅₀ in Rabbits: [Vitavax®]. (Unpublished study received Dec 14 1966 under 400-EX-27; prepared by Food and Drug Research Laboratories, Inc. for Naugatuck Chemical, Div. of United States Rubber Co., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:123429-O)
00003065	Carson, S. (1965) Approximate Acute Oral LD ₅₀ in Rats: [Vitavax®]. (Unpublished study received Dec 14, 1966 under 400-EX-27; prepared by Food and Drug Research Laboratories, Inc., for Naugatuck Chemical, Div. of United States Rubber Co., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:123429-N)
00002879	Chevron Chemical Company (19??) Product Chemistry Data for Difolatan Vitavax®1-2 Flowable Seed Protectant. (Unpublished study received Apr 20, 1976 under 239-2448; CDL:224901-A)
00003321	Chevron Chemical Company (19??) Product Chemistry Data for Difolatan Vitavax®2-2 Flowable Seed Protectant. (Unpublished study including letter dated Aug 15, 1977 from J. Abell to L.R. Stelzer, received Jul 6, 1977 under 239-2459; CDL:230830-A)
00002893	Chevron Chemical Company (19??) Product Chemistry Data for Orthocide Vitavax®1-2 Seed Protectant. (Unpublished study received Apr 23, 1976 under 239-2451; CDL:224838-A)
00003024	Chevron Chemical Company (19??) Product Chemistry Data for Orthocide Vitavax®20-20 Seed Protectant. (Unpublished study received May 11, 1977 under 239-2458; CDL:230221-A)

MRID	CITATION
00003026	Chevron Chemical Company (1977) Summary: Toxicology Data for Orthoicide-Vitavax®20-20 Seed Protectant. Summary of studies 230224-B through 230224-G. (Unpublished study received May 11, 1977 under 239-2458; CDL:230224-A)
00002877	Chevron Chemical Company (1977) Summary: 07--Toxicology Data for Ortho Difolatan-Vitavax®2-2 Flowable Seed Protectant. Summary of studies 230827-B through 230827-G. (Unpublished study received Jul 6, 1977 under 239-2459; CDL:230827-A)
00002884	Chevron Chemical Company (1975) Summary of Toxicology Data for Difolatan Vitavax®1-2 Flowable Seed Protectant. Summary of studies 230799-B through 230799-F. (Unpublished study received Jul 22, 1975 under 239-2448; CDL:230799-A)
00003298	Chevron Chemical Company (1975) Summary of Toxicology Data for Orthocide Vitavax®25-50 Seed Protectant. Summary of studies 224836-B through 224836-F. (Unpublished study received Jul 22, 1975 under 239-2451; CDL:224836-A)
00002897	Chevron Chemical Company (1976) AM-1-1024: Captan-Vitavax®by GLC. Method AM-1-1024 dated May 20, 1976. (Unpublished study received on unknown date under 239-2451; CDL:224838-E)
00002894	Chevron Chemical Company (1976) Shelf-Life Test: Orthocide Vitavax®25-50 S.P. (Unpublished study received on unknown date under 239-2451; CDL:224838-B)
00003022	Chevron Chemical Company (1977) Residue Chemistry Data to Support the Label Registration of Difolatan-Vitavax®2-2 Flowable Seed Protectant. Includes method RM-6E-1 dated Aug 18, 1977 entitled: Determination of captafol residues by gas chromatography. (Unpublished study received Jul 6, 1977 under 239-2459; CDL:230829-A)
00003025	Chevron Chemical Company (1977) Residue Chemistry Data to Support the Label Registration of Orthocide-Vitavax®20-20 Seed Protectant. Includes method RM-IF-1 dated Jun 25, 1976 entitled: Determination of Captan residues in crops. (Unpublished study received May 11, 1977 under 239-2458; CDL:230222-A)
00002935	Chin, W.T.; Dannals, L.E.; Kucharczyk, N. (1972) Environmental Fate Studies on Vitavax®: Status Report on PR 70-15. (Unpublished study received Mar 6, 1972 under 2F1191; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:093515-A)
00002929	Chin, W.T.; Dannals, L.E.; Stone, G.M.; Kucharczyk, N. (1972) 14C Residues in Cottonseed Grown from Seed Treated with Vitavax®200 and Vitavax®300 Seed Protectant Containing Vitavax®14C (Phenyl or Hetero Tag): Project No. 7121. (Unpublished study received on unknown date under 0F0939; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:094582-C)

MRID	CITATION
00002941	Chin, W.T.; Kucharczyk, N.; Smith, A.E. (1972) Nature of Vitavax [®] -Derived Bound Residues in Plants. (Unpublished study received Oct 9, 1973 under 3F1318; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:092254-F)
05002795	Chin, W.T.; Kucharczyk, N.; Smith, A.E. (1973) Nature of carboxin (Vitavax [®])-derived bound residues in barley plants. Journal of Agricultural and Food Chemistry 21(3):506-507.
00003042	Chin, W.T.; Stone, G.M.; Smith, A.E. (1969) Fate of D735 in Dogs. (Unpublished study received Jun 14, 1969 under 9G0819; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091420-U)
00003041	Chin, W.T.; Stone, G.M.; Smith, A.E. (1969) Fate of D735 in Soil. (Unpublished study received Jun 14, 1969 under 9G0819; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091420-T)
00003040	Chin, W.T.; Stone, G.M.; Smith, A.E. (1969) Fate of D735 in Water. Includes undated method. (Unpublished study received Jun 14, 1969 under 9G0819; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091420-S)
00003043	Chin, W.T.; Stone, G.M.; Smith, A.E. (1969) Fate of D735 in Wheat and Barley Plants. (Unpublished study received Jun 14, 1969 under 9G0819; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091420-V)
00003044	Chin, W.T.; Stone, G.M.; Smith, A.E. (1969) 14C-Vitavax Studies on Barley, Wheat and Cotton Plants. (Unpublished study received Jun 14, 1969 under 9G0819; Uniroyal Chemical, Bethany, Conn.; CDL:091420-W)
05002176	Chin, W.T.; Stone, G.M.; Smith, A.E. (1970) Degradation of carboxin (Vitavax [®]) in water and soil. Journal of Agricultural and Food Chemistry 18(4):731-732.
05002177	Chin, W.T.; Stone, G.M.; Smith, A.E. (1970) Metabolism of carboxin (Vitavax [®]) by barley and wheat plants. Journal of Agricultural and Food Chemistry 18(4):709-712.
05011382	Chin, W.T.; Stone, G.M.; Smith, A.E. (1971) The fate of carboxin in soil, plants and animals. Pages 271-279, In Pesticide Terminal Residues, Invited Papers from the International Symposium; Feb 17-19, Tel Aviv, Israel. Edited by A.S. Tahori. London, England: Butterworth.
00025478	Chin, W.T.; Stone, G.M.; Smith, A.E.; et al. (1969) Fate of Carboxin in soil, plants and animals. Proceedings of the 5th Br. Insectic. Fungic. Conference ? : ? - ? . (Also in unpublished submission received Jan 12, 1970 under 0F0939; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091604-Q)

MRID	CITATION
05004996	Chin, W.T.; Stone, G.M.; Smith, A.E.; Schmeling, B. von (1970) Fate of carboxin in soil, plants and animals. Pages 322-327, In <u>Proceedings of the Fifth British Insecticide and Fungicide Conference</u> ; Nov 17-20, 1969, Brighton, England. Vol. 2. Brighton, England: British Insecticide and Fungicide Conference.
05010705	Chopra, S.L.; Chopra, B. (1972) Effect of Vitavax [®] (5,6-dihydro-3-carboxanilido-2-methyl-1,4 oxathiin (systemic fungicide)) on some metabolic activities in wheat. <u>Indian Journal of Agricultural Chemistry</u> 5(2):69-76.
05010593	Chopra, S.L.; Sethi, R.K.; Sikka, V.K. (1974) Chemical composition of chick peas (<u>Cicer arietinum</u>) as affected by Vitavax (5,6-dihydro-3-carboxanilido-2-methyl-1,4-oxathiin). <u>Indian Journal of Agricultural Chemistry</u> 7(1):7-13.
05002888	Coles, C.J.; Singer, T.P.; White, G.A.; Thorn, G.D. (1978) Studies on the binding of carboxin analogs to succinate dehydrogenase. <u>Journal of Biological Chemistry</u> 253(16):5573-5578.
05003675	Clark, R.V. (1971) Influence of Vitavax seed treatment and loose smut infection on yields of barley and wheat. <u>Canadian Plant Disease Survey</u> 51(3):131-133.
00003300	Collier, Y.J.; Womer, J.M.; Stone, G.M.; Kucharczyk, N. (1974) Data of Vitavax [®] -14C Derived Residues in Peanut Hulls, Seed, Oil and Meal. (Unpublished study received on unknown date under 4F1499; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:093979-E)
05002759	Corbeil, M.A.; Curcummelli-Rodostamo, M.; Fanning, R.J.; Graham, B.A.; Kulka, M.; Pierce, J.B. (1973) Ring opening of dihydro-1,4-oxathiins. <u>Canadian Journal of Chemistry</u> 51(16):2650-2658.
05003946	Cors, F.; Kummert, J.; Semal, J. (1971) Some effects of systemic fungicides on virus multiplication in plants. Mededelingen van de Faculteit Landbouwwetenschappen, Rijksuniversiteit Gent. [Communications of the Faculty of Agricultural Sciences, State University of Ghent.] 36(3):1066-1070.
05005447	Couch, R.C.; Dorough, H.W. (1977) Comparative metabolism of selected fungicides. Pages 127-157, In <u>Fate of Pesticides in Large Animals</u> . Edited by G.W. Ivie and H. Dorough. New York: Academic Press.
05003947	Curley, R.L.; Burton, J.C. (1975) Compatibility of <u>Rhizobium japonicum</u> with chemical seed protectants. <u>Agronomy Journal</u> 67(6):807-808.

MRID	CITATION
00003114	Dannals, L.E.; Campbell, C.R.; Cardona, R.A. (1976) Environmental Fate Studies on Vitavax®: Status Report II on PR 70-15. Includes three undated methods. (Unpublished study received Mar 17, 1976 under 400-80; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:223866-A)
05004277	Davis, R.A.; Grahame, R.E.; Kulka, M., inventors; Uniroyal, assignee (1972) Control of virus diseases of plants with carboxamido oxathiins. U.S. patent 3,657,449. Apr 18. 4 p. Int. Cl. A Oln 9/12; U.S. Cl. 424-276.
05002885	Day, D.A.; Arron, G.P.; Laties, G.G. (1978) The effect of carboxins on higher plant mitochondria. FEBS Letters 85(1):99-102.
05005076	Day, E.W., Jr.; Koons, J.R.; Ballee, D.L.; Duane, W.C.; Stallard, D.E.; Wolfe, A.L.; Lynch, V.P.; Edwards, M.J.; Wood, J.S., Jr.; Nishi, K.; Konishi, K.; Stone, G.M. (1976) Fungicides. Pages 251-261, 263-283, 285-297, 299-331, In Analytical Methods for Pesticides and Plant Growth Regulators. Vol. 8: Government Regulations, Pheromone Analysis, Additional Pesticides. New York: Academic Press.
05005466	Dietz, H.J.; Lehmann, H.; Knoll, F.; Grunzel, H., inventors; Veb Fahlberg-List Magdeburg Chemische und Pharmazeutische Fabriken, assignee (1975) Seed disinfecting means. British patent specification 1395449. May 29. 4 p. Int. Cl. 2? A Oln 9/02 21/00 9/12 9/20 9/30.
00003225	Dzialo, D.G.; Lacadie, J.A. (1978) Aerobic Soil Study of 14C-Vitavax® in Sandy Soil: Project No. 7746-1. (Unpublished study received Dec 27, 1978 under 400-80; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:236662-F)
00003226	Dzialo, D.G.; Lacadie, J.A.; Cardona, R.A. (1978) Anaerobic Soil Metabolism of 14C-Vitavax® in Sandy Soil: Project No. 7751. (Unpublished study received Dec 27, 1978 under 400-80; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:236662-G)
05012997	Eade, R.P. (1972) Effect of carboxin on loose smut of wheat. Pages 155-159, In Proceedings of the New Zealand Weed and Pest Control Conference. Vol. 25. Hamilton, New Zealand: New Zealand Weed and Pest Control Conference.
00003060	Edgington, L.V. (19??) Bioassay of 735. Undated method. (Unpublished study received Dec 14, 1966 under 400-EX-27; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:123429-C)
05002835	El-Dib, M.A.; Aly, O.A. (1972) Colorimetric determination of phethylamide pesticides in natural waters. Journal of the Association of Official Analytical Chemists 55(6):1276-1279.

MRID	CITATION
05006154	El-Dib, M.A.; Aly, O.A. (1976) Persistence of some phenylamide pesticides in the aquatic environment--I. Hydrolysis. Water Research 10(12):1047-1050.
05003915	El-Dib, M.A.; Aly, O.A. (1976) Persistence of some phenylamide pesticides in the aquatic environment--I. Hydrolysis--II. Adsorption on clay minerals--III. Biological degradation. Water Research 10(12):1047-1053,1055-1059.
05003218	El-Dib, M.A.; Aly, O.A. (1976) Persistence of some phenylamide pesticides in the aquatic environment--III. Biological degradation. Water Research 10(12):1055-1059.
05002882	El-Dib, M.A.; Aly, O.A. (1977) Removal of phenylamide pesticides from drinking waters--I. Effect of chemical coagulation and oxidants. Water Research 11(8):611-616.
05002884	El-Dib, M.A.; Aly, O.A. (1977) Removal of phenylamide pesticides from drinking waters--II. Adsorption on powdered carbon. Water Research 11(8):617-620.
05007883	Engelhardt, G.; Wallnoefer, P. (1976) Hydrolyse von methoxysubstituierten Phenylharnstoffen, Acylaniliden und Phenylcarbamaten durch eine mikrobielle Aryl Acylamidase [Hydrolysis of methoxysubstituted phenylureas, acylanilides and phenylcarbamates by a microbial aryl acylamidase.] Zentralblatt fuer Bakteriologie, Parasitenkunde, Infektionskrankheiten und Hygiene, Abteilung 1: Originale, Reihe B 162(2):138-144.
05002883	Engelhardt, G.; Wallnoefer, P.R.; Plapp, R. (1971) Degradation of linuron and some other herbicides and fungicides by a linuron-inducible enzyme obtained from <u>Bacillus sphaericus</u> . Applied Microbiology 22(3):284-288.
05002833	Farrow, J.E.; Hoodless, R.A.; Hopkinson, A. (1975) Fungicide residues. Part IV. Determination of residues of carboxin in grain by gas chromatography. Analyst 100(1189):249-252.
00003139	Fink, R. (1974) Final Report: Eight-Day Dietary LC ₅₀ --Mallard Ducks: Project No. 117-102. (Unpublished study including official analytical report, received Mar 11, 1974 under 400-106; prepared by Truslow Farms, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:128723-A)
05002757	Fisher, D.J. (1976) Effects of some fungicides on <u>Rhizobium trifolii</u> and its symbiotic relationship with white clover. Pesticide Science 7(1):10-18.
00025477	Frederiksen, R.A.; Reyes, L. (1969) Seed Treatments, Seedling Stands and Seedling Disease. (Unpublished study received Jan 12, 1970 under OF0939; prepared by Texas A & M Univ., Agricultural Research Station, Dept. of Plant Sciences, submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091604-P)

MRID	CITATION
05002830	Garg, F.C. (1977) Effect of systemic fungicides on the growth of <u>Azotobacter</u> spp. <i>Pesticides</i> 11(4):41.
05006926	Garg, F.C.; Tauro, P.; Grover, R.K. (1973) Effect of fungitoxicants on <u>Rhizobia</u> . <i>Indian Journal of Microbiology</i> 13(3):179-181.
00002994	Geise, T.E. (19??) Vitavax [®] HB-25 Production Procedure. (Unpublished study received Jul 15, 1976 under 400-118; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:24929-B)
05006912	Gross, Y.; Kenneth, R. (1973) The fate of some systemic fungicides in plants when taken up by way of soil. <i>Phytoparasitica</i> 1(1):60.
00025469	Hansing, E.D. (1969) [Efficacy Data for Vitavax [®] on Wheat]. Prelim. rept. (Unpublished study received Jan 12, 1970 under 0F0939; prepared by Kansas State Univ., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091604-H)
00025474	Hansing, E.D. (1969) [Efficacy Data for Vitavax [®] on Sorghum]. (Unpublished study received Jan 12, 1970 under 0F0939; prepared by Kansas State Univ., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091604-M)
05005474	Hardison, J.R. (1971) Relationships of molecular structure of 1,4-oxathiin fungicides to chemotherapeutic activity against rust and smut fungi in grasses. <i>Phytopathology</i> 61(6):731-735.
05003951	Harri, J.A.; Larsen, P.O.; Powell, C.C., Jr. (1975) Bacterial leaf spot and blight of Rieger begonia cv Aphrodite Rose. Pages 4-6, <u>In</u> Ohio Florists' Association, Bulletin no. 552.
00003310	Hathaway, D. (1970) Report to Uniroyal Chemical, Division of Uniroyal, Inc.: Acute Dust Inhalation Toxicity Study on Vitavax Seed Protectant with Captan: IBT No. N8586. (Unpublished study received May 8, 1972 under 2F1191; prepared by Industrial Bio-Test Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091003-E)
00004244	Hathaway, D. (1970) Report to Uniroyal Chemical, Division of Uniroyal, Inc.: Acute Dust Inhalation Toxicity Study on Vitavax Seed Protectant with Thiram: IBT No. N8586. (Unpublished study received May 8, 1972 under 2F1191; prepared by Industrial Bio-Test Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091003-F)
05003603	Haware, M.P. (1973) Effect of systemic fungicides on <u>Fusarium lini</u> in vitro and their up take by linseed seedlings. <i>Indian Journal of Mycology and Plant Pathology</i> 2(2):136-138.

MRID

CITATION

- 05005919 Hennings, P.; Goehmann, P.; Otterpohl, G. (1976) Zur Reaktion von 1,4-Oxathienen mit Jodwasserstoffsäure und ihre Anwendung zur Bestimmung des Fungizids Carboxin. [Reaction of 1,4-oxathienes with hydroiodic acid and its application to the determination of the fungicide carboxin.] Zeitschrift fuer Chemie 16(11):448-449.
- 05004445 Hey, G.L., inventor; Murphy Chemical Limited, assignee (1973) Improvements in fungicides. British patent specification 1,337,454. Nov 14. 4 p. Int. Cl. A 01N 9/02,9/12,9/20,21/00.
- 05003787 Hill, K.R. (1975) IUPAC Commission on Terminal Residues. Journal of the Association of Official Analytical Chemists 58(6):1256-1285.
- 00003035 Holsing, G.C. (1968) Final Report: Acute Eye Irritation--Rabbits: Project No. 798-151. (Unpublished study received Jun 14, 1969 under 9G0819; prepared by Hazleton Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091420-L)
- 00021625 Holsing, G.C. (1968) Final Report: Acute Eye Irritation--Rabbits: Project No. 798-151. (Unpublished study received Jul 19, 1968 under 400-80; prepared by Hazleton Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:003258-G)
- 00003216 Holsing, G.C. (1968) Final Report: Repeated Dermal Application--Rabbits: Project No. 798-148. (Unpublished study received Jan 13, 1969 under 400-EX-33; prepared by Hazleton Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL: 123439-M)
- 00003032 Holsing, G.C. (1968) Final Report: Three-Generation Reproduction Study--Rats: Project No. 798-104. (Unpublished study received Jun 14, 1969 under 9G0819; prepared by Hazleton Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL: 091420-G)
- 00021626 Holsing, G.C. (1968) Summary: Repeated Dermal (Leary Design)--Rabbits: Project No. 798-148. (Unpublished study received Jul 19, 1968 under 400-80; prepared by Hazleton Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:003258-K)
- 00003030 Holsing, G.C. (1969) Final Report: Two-Year Dietary Administration--Dogs: Project No. 798-103. (Unpublished study received Jun 14, 1969 under 9G0819; prepared by Hazleton Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL: 091420-E)
- 00003031 Holsing, G.C. (1969) Final Report: 24-Month Dietary Administration--Albino Rats: Project No. 798-102. (Unpublished study received Jun 14, 1969 under 9G0819; prepared by Hazleton Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091420-F)

MRID	CITATION
00003152	Holsing, G.C. (1969) Final Report: 24-Month Dietary Administration--Albino Rats: Project No. 798-102. (Unpublished study received Oct 21, 1971 under 400-90; prepared by Hazleton Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:050926-A)
00002914	Horne, C.W.; Philley, G.L.; Smith, L.R. (1972) Results of 1972 Extension Peanut Disease Control Demonstrations Conducted at Mason, Grapeland, Dublin, and Floresville, Texas. (Unpublished study received May 1, 1974 under 4F1499; prepared by Texas A & M Univ., Texas Agricultural Extension Service, submitted by Uniroyal Chemical, Bethany, Conn.; CDL:094551-U)
00003010	Imlay, P. (1977) Report: The Acute Dermal LD50 of Vitavax®200 Flowable Fungicide on New Zealand Albino Rabbits. (Unpublished study received Jul 14, 1978 under 400-112; prepared by Cannon Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:235042-B)
05006081	Karchik, O.N.; Loginova, L.N.; Eglit, I.E. (1975) Deistvie proizvodnykh oksatiina na kataliticheskuyu aktivnost' metallosoderzhashchikh oksidoreduktaz pshenitsy. [Effect of oxathiin derivatives on the catalytic activity of metal-containing wheat oxidoreductases.] Khimiya v Sel'skom Khozyaistve. [Chemistry in Agriculture.] 13(2):119-121.
00002943	Kennedy, G. (1971) Report to Uniroyal Chemical, Division of Uniroyal, Inc.: Distribution and Excretion Study with Vitavax in Albino Rats: IBT No. E9289. (Unpublished study received Oct 9, 1973 under 3F1318; prepared by Industrial Bio-Test Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:092254-H)
00002944	Kennedy, G. (1971) Report to Uniroyal Chemical, Division of Uniroyal, Inc.: Characterization of 14C in Excreta of Rats Following Oral Administration of Vitavax: IBT No. E9290. (Unpublished study received Oct 9, 1973 under 3F1318; prepared by Industrial Bio-Test Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:092254-I)
00021650	Kennedy, G.; Jenkins, D.H. (1971) Report to Uniroyal Chemical, Division of Uniroyal, Inc.: Milk and Meat Residue Study in Dairy Cows Treated with 14C-Labeled Vitavax®: IBT No. J294. (Unpublished study received Jul 16, 1979 under 3F1318; prepared by Industrial Bio-Test Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:098393-B)
00002945	Kennedy, G.; Jenkins, D.H. (1971) Report to Uniroyal Chemical, Division of Uniroyal, Inc.: Milk and Meat Residue Study in Dairy Cows Treated with 14C-Labeled Vitavax®: IBT No. J294. (Unpublished study received Oct 9, 1973 under 3F1318; prepared by Industrial Bio-Test Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:092254-J)

MRID	CITATION
05003664	Kirk, B.T.; Sinclair, J.B.; Lambremont, E.N. (1969) Translocation of 14C-labeled chloroneb and IMOC in cotton seedlings. <i>Phytopathology</i> (59):1473-1476.
00003120	Knickerbocker, M. (1977) Teratologic Evaluation of Vitavax® Technical in Sprague Dawley Rats. (Unpublished study recieved Jan 24, 1978 under 400-81; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:233500-F)
05005992	Knoll, F.A. (1978) Beeinflussung eines gegen <u>Verticillium albo-atrum</u> Rke. et Berth. wirksamen phyto-genen Hemmstoffpotentials beim Einsatz und Bionachweis von Systemfungiziden. [Influence of a phytogenic inhibitor potential effective against <u>Verticillium albo-atrum</u> Rke. et Berth. in the use and bio-assay of systemic fungicides.] <i>Archiv fuer Phytopathologie und Pflanzenschutz</i> 14(1):17-26.
05013365	Kollmorgen, J.F.; Ballinger, D.J. (1975) Suppression of the emergence of wheat and barley by mancozeb. <i>Australian Journal of Experimental Agriculture and Animal Husbandry</i> 15(76):700-704.
05004125	Kommedahl, T.; Lang, D.S. (1973) Effect of temperature and fungicides on survival of corn grown from kernals infected with <u>Helminthosporium maydis</u> . <i>Phytopathology</i> 63(1):138-140.
05009410	Konashevich, V.A.; Katseval, A.M. (1972) Preparaty protiv pyl'noy golovni pshenitsy. [Control agents for loose smut of wheat.] <i>Zashchita Rastenii</i> (Moscow). [Plant Protection.] 17(2):35.
05003529	Kratka, J. (1975) Some biochemical changes in young barley plants, due to the Vitavax disinfection of seeds against <u>Ustilago nuda</u> (Jens.) Rostr. <i>Zentralblatt fuer Bakteriologie, Parasitenkunde, Infektionskrankheiten und Hygiene, Abteilung II</i> 130(2):165-170.
05007643	Kratka, J.; Prochazka, M.; Ujevic, I. (1978) Vliv fungicidu na nekte-re metabolicke procesy a sladovnickou kvalitu jecmene. [Effect of fungicides on some metabolic processes and on the malting quality of barley.] <i>Agrochemia</i> 18(2):57-60.
05003530	Kratka, J.; Ujevic, I. (1971) The effect of seed disinfection with systemic preparations Vitavax® and Plantvax® on respiration activity and on catalase in barley. <i>Zentralblatt fuer Bakteriologie, Parasitenkunde, Infektionskrankheiten und Hygiene, Abteilung II</i> 126(2):154-161.
05002989	Kritzman, G.; Chet, I.; Henis, Y. (1977) Effect of carbon dioxide on growth and carbohydrate metabolism in <u>Sclerotium rolfsii</u> . <i>Journal of General Microbiology</i> 100(1):167-175.

MRID	CITATION
00005545	Kuc, W.J.; Doebbler, G.F. (1979) Uniroyal Chemical 14C-Vitavax® Bioconcentration Study: Channel Catfish, <i>Ictalurus punctatus</i> (Rafinesque): Project No. 11506-82. Includes undated method. (Unpublished study including metabolite study, received Mar 28, 1979 under 400-81; prepared by Union Carbide Corp., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:098029-G)
00005544	Kuc, W.J.; Doebbler, G.F. (1979) Uniroyal Chemical 14C-Vitavax® Bluegill Sunfish Bioconcentration Study: UCES Project No. 11506-82. Includes undated method. (Unpublished study including metabolite study, received Mar 28, 1979 under 400-81; prepared by Union Carbide Corp., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:098029-F)
00003049	Kucharczyk, N. (1971) Stability of Vitavax® on Treated Seed (Cotton). (Unpublished study received on unknown date under 0F0939; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:093245-I)
05004135	Kuiper, J. (1968) Experiments on the control of cereal smuts with a derivative of 1,4-oxathiin. Australian Journal of Experimental Agriculture and Animal Husbandry 8(31):232-237.
00003228	Lacadie, J.A.; Dannals, L.E. (1978) Vitavax®-14C Laboratory Column Leaching in Sandy Soil: Project No. 7831. (Unpublished study received Dec 27, 1978 under 400-80; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:236662-I)
00003229	Lacadie, J.A.; Dzialo, D.G.; Dannals, L.E. (1978) Vitavax® 14C Aged Laboratory Column Leaching in Sandy Soil: Project No. 7832. (Unpublished study received Dec 27, 1978 under 400-80; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:236662-J)
00003227	Lacadie, J.A.; Gerecke, D.R.; Cardona, R.A. (1978) Vitavax®-14C Laboratory Column Leaching Study in Clay Loam: Project No. 7758. (Unpublished study received Dec 27, 1978 under 400-80; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:236662-H)
00003058	Lane, J.R. (1966) Residue Analysis for D-735 and/or F-461. Method dated May 16, 1966. (Unpublished study received Jan 25, 1967 under 400-81; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:121531-A)
00003183	Lane, J.R. (1968) Residue Analysis for Vitavax®. Method dated Nov 1, 1968. (Unpublished study received Mar 25, 1969 under 9G0819; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:093520-AA)
00026087	Lane, J.R. (1968) Residue Analysis for Vitavax® and/or Plantvax. Method dated Jan 18, 1968. (Unpublished study received Jul 19, 1968 under 400-80; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:003258-P)

MRID	CITATION
05002737	Lane, J.R. (1970) Colorimetric microdetermination of Vitavax® (5,6 dihydro-2-methyl-1, 4-oxathiin-3-carboxanilide) residues in crops. <i>Journal of Agricultural and Food Chemistry</i> 18(3):409-412.
00002947	Lane, J.R. (1972) Determination of Free and Bound Para-Aminophenol Residues in Feces and Urine. Includes method dated Sep 18, 1972. (Unpublished study received Oct 9, 1973 under 3F1318; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:092254-L)
00025467	Lane, J.R.; Siskin, H.R. (1969) The Effects of FDA Tolerance Pesticides upon the Recovery of Vitavax® Residues from Crop Plants: Cotton, Peanut, Sorghum, Wheat and Barley. (Unpublished study received Jan 12, 1970 under 0F0939; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091604-F)
00025468	Lane, J.R.; Siskin, H.R. (1970) Specificity of the Vitavax® Method for Seeds. (Unpublished study received Jan 12, 1970 under 0F0939; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091604-G)
00002958	Leroux, P.; Gredt, M. (1977) Fate of Carboxin Used in Treatment of Wheat Seed: Absorption, Transport and Degradation. (Unpublished study received Mar 6, 1972 under 2F1191; prepared by National Agricultural Research Center, Plant Treatment Laboratory--C.N.R.A. [France], submitted by Uniroyal Chemical, Bethany, Conn.; CDL:093515-C)
05006363	Leroux, P.; Gredt, M. (1972) Devenir de la carboxine utilisee en enrobage de semences de ble: absorption, transport et degradation. [The fate of carboxin used as a dressing for wheat seed: absorption, translocation and degradation.] <i>Phytiatrie-Phytopharmacie</i> 21(1):45-60.
00003033	Littlefield, N.A. (1968) Acute Inhalation Exposure--Rats: Project No. 798-153. (Unpublished study received Jun 14, 1969 under 9G0819; prepared by Hazleton Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091420-J)
00021624	Littlefield, N.A. (1968) Acute Inhalation Exposure--Rats: Project No. 798-153. (Unpublished study received Jul 19, 1968 under 400-80; prepared by Hazleton Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:003258-F)
05002745	Littlejohns, D.A.; McLaren, A.D.; Aylesworth, J.W. (1976) The effect of foliar sprays in controlling ozone damage in white beans. <i>Canadian Journal of Plant Science</i> 56(2):430.
05011365	Luisi, N. (1977) Prodotti sistemici in patologia vegetale, ecologia e adattamento dei patogeni. [Systemic products used in vegetable pathology, ecology and resistance of the pathogens.] <i>Securitas. [Safety.]</i> 62(11/12):659-666.

MRID	CITATION
05003852	Lyr, H.; Ritter, G.; Banasiak, L. (1974) Detoxication of carboxin. Zeitschrift fuer Allgemeine Mikrobiologie 14(4):313-320.
05012882	Macias, W.; Smoter, J. (1973) Badania nad fitotoksycznoscia oraz wplywem fungicydow z grupy PCNB i TMTD na stan odzywienia cebuli. [Phytotoxic effects of fungicides of the PCNB and TMTD groups on the nourishment of onions.] Biuletyn Warzywniczy. [Bulletin of Vegetable Crops Research Work.] XV:249-268.
000025484	Martinson, C.A.; Neidermyer, R.W.; Rouwenhorst, D.; et al. (1973) Reasoning in Support of the Petition. (Unpublished study received Jun 17, 1974 under 5F1525; prepared in cooperation with Iowa State Univ. and Univ. of Illinois, submitted by Uniroyal Chemical, Bethany, Conn.; CDL:094043-I)
00003309	Mastalski, K. (1970) Report to Uniroyal Chemical, Division of Uniroyal, Inc.: Acute Oral Toxicity Study on Vitavax®Seed Protectant with Captan in Bobwhite Quail: IBT No. J8584. (Unpublished study received May 8, 1972 under 2F1191; prepared by Industrial Bio-Test Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091003-D)
00003308	Mastalski, K. (1970) Report to Uniroyal Chemical, Division of Uniroyal, Inc.: Acute Oral Toxicity Study on Vitavax®Seed Protectant with Thiram in Bobwhite Quail: IBT No. J8584. (Unpublished study received May 8, 1972 under 2F1191; prepared by Industrial Bio-Test Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091003-C)
00002923	Mastri, C. (1970) Report to Uniroyal Chemical, Division of Uniroyal, Inc.: Four-Day Fish Toxicity Study on Three Samples of Vitavax®Seed Protectants: IBT No. A8585. (Unpublished study received on August 8, 1975 under 400-118; prepared by Industrial Bio-Test Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:230400-G)

MRID	CITATION
00021610	Mastri, C. (1970) Report to Uniroyal Chemical, Division of Uniroyal, Inc.: Four-Day Fish Toxicity Study on Three Samples of Vitavax [®] Seed Protectants: IBT No. A8585. (Unpublished study received Feb 18, 1972 under 400-81; prepared by Industrial Bio-Test Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:003261-D)
05004346	Mathre, D.E. (1971) Mode of action of oxathiin systemic fungicides: III. Effect on mitochondrial activities. Pesticide Biochemistry and Physiology 1(2):216-224.
05002488	Mathre, D.E. (1972) Effects of oxathiin systemic fungicides on various biological systems. Bulletin of Environmental Contamination and Toxicology 8(5):311-316.
05004904	Mathre, D.E.; Stewart, V.R.; Johnston, R.H.; Baldrige, D.E. (1975) Fungicide treatment of wheat seed: is it necessary? Journal of Environmental Quality 4(1):117-120.
00003316	Matthews, R.J. (1970) Acute Dermal Application--Rabbits: Vitavax [®] with Captan: Final Report. (Unpublished study received May 8, 1972 under 2F1191; prepared by Pharmakon Laboratories, submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091003-L)
00003315	Matthews, R.J. (1970) Acute Dermal Application--Rabbits: Vitavax [®] with Thiram: Final Report. (Unpublished study received May 8, 1972 under 2F1191; prepared by Pharmakon Laboratories, submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091003-K)
00003314	Matthews, R.J. (1970) Acute Dermal Application--Rabbits: Vitavax [®] : Final Report. (Unpublished study received May 8, 1972 under 2F1191; prepared by Pharmakon Laboratories, submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091003-J)
00003319	Matthews, R.J. (1970) Acute LD ₅₀ --Rats (Oral): Vitavax [®] with Captan: Final Report. (Unpublished study received May 8, 1972 under 2F1191; prepared by Pharmakon Laboratories, submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091003-O)
00003318	Matthews, R.J. (1970) Acute LD ₅₀ --Rats (Oral): Vitavax [®] with Thiram: Final Report. (Unpublished study received May 8, 1972 under 2F1191; prepared by Pharmakon Laboratories, submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091003-N)
00003317	Matthews, R.J. (1970) Acute LD ₅₀ --Rats (Oral): Vitavax [®] : Final Report. (Unpublished study received May 8, 1972 under 2F1191; prepared by Pharmakon Laboratories, submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091003-M)
00003328	Matthews, R.J. (1970) Draize Eye Irritation--Rabbits: Vitavax [®] : Final Report. (Unpublished study received Feb 18, 1972 under 400-81; prepared by Pharmakon Laboratories, submitted by Uniroyal Chemical, Bethany, Conn.; CDL:003260-A)

MRID	CITATION
00003329	Matthews, R.J. (1970) Draize Eye Irritation--Rabbits: Vitavax [®] with Captan: Final Report. (Unpublished study received Feb 18, 1972 under 400-81; prepared by Pharmakon Laboratories, submitted by Uniroyal Chemical, Bethany, Conn.; CDL:003260-B)
00003313	Matthews, R.J. (1970) Primary Skin Irritation--Rabbits: Vitavax [®] with Captan: Final Report. (Unpublished study received May 8, 1972 under 2F1191; prepared by Pharmakon Laboratories, submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091003-I)
00003312	Matthews, R.J. (1970) Primary Skin Irritation--Rabbits: Vitavax [®] with Thiram: Final Report. (Unpublished study received May 8, 1972 under 2F1191; prepared by Pharmakon Laboratories, submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091003-H)
00003311	Matthews, R.J. (1970) Primary Skin Irritation--Rabbits: Vitavax [®] : Final Report. (Unpublished study received May 8, 1972 under 2F1191; prepared by Pharmakon Laboratories, submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091003-G)
00003164	Matthews, R.J. (1970) Primary Skin Irritation--Rabbits: Vitavax [®] : Final Report. (Unpublished study received Apr 21, 1977 under unknown admin. no.; prepared by Pharmakon Laboratories, submitted by Uniroyal Chemical, Bethany, Conn.; CDL:229370-K)
00003012	Matthews, R.J. (1973) Acute LD ₅₀ --Rats, Oral: Final Report. (Unpublished study received Jul 14, 1978 under 400-112; prepared by Pharmakon Laboratories, submitted by Uniroyal Chemical, Bethany, Conn.; CDL:235042-D)
05004129	Michail, S.H.; Elarosi, H.; Abd-El-Rehim, M.A.; Shohda, W.T. (1975) Persistence of certain systemic and non-systemic fungicides in the soil, with special reference to the method of application. <i>Phytopathologia Mediterranea</i> 14(2/3):138-139.
05002754	Mohan, S.T. (1975) Cytological effects of fungicides, plantvax and vitavax on somatic cells of <u>Allium cepa</u> [letter]. <i>Current Science</i> 44(22):813-814.
00002926	Morse Laboratories, Incorporated (1975) Residues in PPM: [Vitavax [®]]. (Unpublished study received Mar 27, 1975 under 400-118; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:230407-J)
00033019	Morse Laboratories, Incorporated (1980) [Vitavax [®] Residues in Peanut Seeds]. (Unpublished study received Jun 3, 1980 under OK 80/9; prepared in cooperation with Uniroyal Chemical, submitted by Oklahoma, Dept. of Agriculture, Oklahoma City, Okla.; CDL:242612-D)

MRID

CITATION

- 05012734 Mueller, W.; Schewe, T. (1976) Zur Wirkung einiger Atmungshemmstoffe auf den Tetramethyl-p-phenylenediamin-Kurzschlussweg in nicht-phosphorylierenden Rinderherz-Elektronentransportpartikeln. [Action of some respiratory inhibiting substances on the tetramethyl-p-phenylenediamine bypass in nonphosphorylating bovine heart electron transport particles.] *Acta Biologica et Medica Germanica* 35(12):1737-1740.
- 05019626 Mueller, W.; Schewe, T.; Lyr, H.; Zanke, D. (1977) Wirkmechanismus der Atmungshemmung durch die Systemfungicide der Carboxingruppe Wirkung von Oxathiinderivaten und -analoge auf nichtphosphorylierende submitochondriale Partikeln aus Rinderherz sowie Trametes versicolor und Trichoderma viride. [Action mechanism of respiratory inhibition by systemic fungicides of the carboxin group. Effects of oxathiin derivatives and analogues on nonphosphorylating submitochondrial particles from bovine heart, Trametes versicolor and Trichoderma viride.] *Zeitschrift fuer Allgemeine Mikrobiologie* 17(5):359-372.
- 05006904 Mukherji, S.K.; Bagchi, B.N. (1977) Efficacy of Vitavax[®]-200 against some fungi in vitro. *Pesticides* 11(11):32-33.
- 05001446 Mukhopadhyay, A.N. (1976) Recent advances in the control of Sclerotium root rot of sugarbeet. *Indian Phytopathology* 29(1):102.
- 05001444 Mukhopadhyay, A.N.; Tewari, K.C. (1975) Control of sclerotium root rot of sugar beet. *Indian Phytopathology* XXVIII(1):140.
- 05001203 Mukhopadhyay, A.N.; Thakur, R.P. (1971) Control of Sclerotium root rot of sugarbeet with systemic fungicides. *Plant Disease Reporter* 55(7):630-634.
- 05002341 Mulholland, R.I. (1972) Bunt control in wheat. Pages 165-168, In *Proceedings of the New Zealand Weed and Pest Control Conference*. Vol. 25. Hamilton, N.Z.: Ruakura Agricultural Research Centre.
- 05004588 Mulholland, R.I. (1972) Loose smut control in wheat. Pages 152-154, In *Proceedings of the New Zealand Weed and Pest Control Conference*. Vol. 25. Hamilton, N.Z.: New Zealand Weed and Pest Control Conference.
- 05004589 Mulholland, R.I. (1976) Control of loose smut of oats. Pages 217-220, In *Proceedings of the 29th New Zealand Weed and Pest Control Conference*. Hamilton, N.Z.: New Zealand Weed and Pest Control Conference.
- 05006573 Muller, B. (1968) Qu'est-ce que la carboxine? [What is carboxin?] *Defense des Vegetaux* 22(133):239-248.

MRID	CITATION
05011379	Murdoch, C.L.; Lacmlen, F.F.; Parvin, P.E. (1973) Fungicidal control of <u>Puccinia striiformis</u> on three cultivars of <u>Poa pratensis</u> in Hawaii. <u>Plant Disease Reporter</u> 57(3):217-219.
05010986	Mustafa, M.; D'yakov, Y.T. (1979) Vliyanie pestitsidov na porazhaemost' kartofelya i tomatov fitoftorozom. [Effect of pesticides on the susceptibility of potatoes and tomatoes to phytophthorosis.] <u>Khimiya v Sel'skom Khozyaistve</u> . [Chemistry in Agriculture.] 17(1):56-59.
05001234	Muthusamy, S. (1973) Systemic fungicides in the control of pineapple disease of sugar cane. <u>Sugarcane Pathologists' Newsletter</u> 10:14-15.
05010704	Nayak, M.L.; Verma, J.P. (1975) Uptake and translocation of chemicals in cotton seedlings. <u>Zeitschrift fuer Pflanzenkrankheiten und Pflanzenschutz</u> 82(10):587-592.
00003070	Nichols, D.F. (1967) Monthly Report: [Fungicides: Vitavax® and Plantvax®.] (Unpublished study received Dec 6, 1967 under 400-EX-30; submitted by Uniroyal Chemical, Bethany, Conn.; CDL: 123434-G)
05005989	Nilova, V.P.; Ksendzova, E.N. (1972) Vliyanie vitavaksa na aktivnost' katalazy i vospriimchivost' pshenitsy k buroi rzhavchine. [Effects of vitavax on catalase activity and susceptibility of wheat plants to brown rust.] <u>Mikologiya i Fitopatologiya</u> . [Mycology and Phytopathology.] 6(4):355-358.
05002761	Onuska, F.I.; Comba, M.E. (1974) Sodium azide-iodine reagent for the detection of 1,4-oxathiin derivatives by thin-layer chromatography. <u>Journal of Chromatography</u> 100(1):247-248.
05002893	Onuska, F.I.; Comba, M.E.; Harrison, A.G. (1976) Chemical ionization and high resolution electron impact mass spectra of 5,6-dihydro-2-methyl-1,4-oxathiin-3-carboxanilide and three of its metabolites. <u>Biomedical Mass Spectrometry</u> 3(5):248-254.
00003063	Ozer, B.L. (1966) Report: Subacute (90-day) Feeding Studies with D-735 in Rats. (Unpublished study received Dec 14, 1966 under 400-EX-27; prepared by Food and Drug Research Laboratories, Inc. for United States Rubber Co., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:123429-F)
05012255	Padalkin, I.I. (1978) Toksichnost' vitavaksa dlia kur. [Toxicity of vitavax in hens.] <u>Veterinariya (Moscow)</u> . [Veterinary Science.] (6):85-86.
05003601	Peterson, C.A.; Edgington, L.V. (1975) Penetration and retention of systemic pesticides in the symplast of potato tuber tissue. Pages 28, In <u>Proceedings of the Canadian Phytopathological Society</u> . Vol. 42.

MRID	CITATION
05006896	Peterson, C.A.; Edgington, L.V. (1976) Entry of pesticides into the plant symplast as measured by their loss from an ambient solution. <i>Pesticide Science</i> 7(5):483-491.
05002887	Plato, C. (1972) Differential scanning calorimetry as a general method for determining purity and heat of fusion of high-purity organic chemicals: application to 64 compounds. <i>Analytical Chemistry</i> 44(8):1531-1534.
00003323	Puchalsky, C.B. (1966) Infrared Methods for the Assay of D-735 and F-461, and Their Content in Formulations. Method dated Apr 6, 1966. (Unpublished study received on January 12, 1970 under 0F0939; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:093245-K)
00003172	Puchalsky, C.B. (1968) Infrared Method for the Assay of D-735, and Its Content in Formulations. Method dated Jul 15, 1968. (Unpublished study received Mar 25, 1969 under 9G0819; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:093520-H)
00024927	Puchalsky, C.B. (1976) Infrared Method for Captan and Vitavax® in UBI-1107 and 1109. Method dated Mar 22, 1976. (Unpublished study received on unknown date under 0F0939; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:098575-B)
00002996	Puchalsky, C.B. (1976) Infrared Method for Captan and Vitavax® in UBI-1107 and 1109. Method dated Mar 22, 1976. (Unpublished study received Jul 15, 1976 under 400-118; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:224929-D)
00005861	Puchalsky, C.B. (1979) Infrared Method for the Vitavax® Content of Vitavax®-4G. Method dated Jan 30, 1979. (Unpublished study received Feb 6, 1979 under 400-EX-58; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:237329-C)
00025476	Ranney, C.D.; Burchfield, E.G. (1969) [Vitavax® Used in Cottonseed Treatment]. (Unpublished study received Jan 12, 1970 under 0F0939; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091604-O)
00025471	Reed, H.E.; Chambers, A.Y.; Richard, G.; et al. (1969) [Efficacy in Barley]. (Unpublished study including published data, received Jan 12, 1970 under 0F0939; prepared in cooperation with Univ. of Tennessee, Agricultural Experiment Station, Dept. of Agricultural Biology and others, submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091604-J)
00025470	Richard, G.; Vallier, J.P. (1969) Treatment of cereal seed by the combination of Carboxin with copper oxyquinolate. <i>Proceedings of the 5th Br. Insectic. Fungic. Conference</i> :45-54. (Also <u>In</u> unpublished submission received Jan 12, 1970 under 0F0939; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091604-I)

MRID

CITATION

- 00003297 Rittenhouse, J.R.; Narcisse, J.K. (1975) The Acute Dermal Toxicity of Difolatan-Vitavax® 1-2F Seed Protectant: S-799: SOCAL 703/XV: 98. (Unpublished study received Jul 22, 1975 under 239-2448; prepared by Standard Oil Co. of California, Safety Div., submitted by Chevron Oil Co., Richmond, Calif.; CDL:230799-C)
- 00003299 Rittenhouse, J.R.; Narcisse, J.K. (1975) The Acute Dermal Toxicity of Orthocide-Vitavax® 25-50 W.P. Seed Protectant: S-805: SOCAL 710/XV:99. (Unpublished study received Jul 22, 1975 under 239-2451; prepared by Standard Oil Co. of California, Safety Div., submitted by Chevron Chemical Co., Richmond, Calif.; CDL: 224836-C)
- 00002892 Rittenhouse, J.R.; Narcisse, J.K. (1975) The Acute Inhalation Toxicity of Orthocide-Vitavax® 25-50 W.P. Seed Protectant: S-808: SOCAL 713/XXI:55. (Unpublished study received Jul 22, 1975 under 239-2451; prepared by Standard Oil Co. of California, Safety Div., submitted by Chevron Chemical Co., Richmond, Calif.; CDL:224836-F)
- 00002889 Rittenhouse, J.R.; Narcisse, J.K. (1975) The Acute Oral Toxicity of Orthocide-Vitavax® 25-50 W.P. Seed Protectant: S-804: SOCAL 709/XVIII:135. (Unpublished study received Jul 22, 1975 under 239-2451; prepared by Standard Oil Co. of California, Safety Div., submitted by Chevron Chemical Co., Richmond, Calif.; CDL: 224836-B)
- 00002885 Rittenhouse, J.R.; Narcisse, J.K. (1975) The Acute Oral Toxicity of Difolatan-Vitavax® 1-2F Seed Protectant: S-798: SOCAL 702/XVIII: 133. (Unpublished study received Jul 22, 1975 under 239-2448; prepared by Standard Oil Co. of California, Safety Div., submitted by Chevron Chemical Co., Richmond, Calif.; CDL:230799-B)
- 00002888 Rittenhouse, J.R.; Narcisse, J.K. (1975) The Acute Vapor Toxicity of Difolatan-Vitavax® 1-2F Seed Protectant: S-802: SOCAL 706/XXI: 48. (Unpublished study received Jul 22, 1975 under 239-2448; prepared by Standard Oil Co. of California, Safety Div., submitted by Chevron Chemical Co., Richmond, Calif.; CDL:230799-F)
- 00002886 Rittenhouse, J.R.; Narcisse, J.K. (1975) The Eye Irritation Potential of Difolatan-Vitavax® 1-2F Seed Protectant (CC 5998): S-800: SOCAL 704/XXII:16. (Unpublished study received Jul 22, 1975 under 239-2448; prepared by Standard Oil Co. of California, Safety Div., submitted by Chevron Chemical Co., Richmond, Calif.; CDL:230799-D)
- 00002890 Rittenhouse, J.R.; Narcisse, J.K. (1975) The Eye Irritation Potential of Orthocide-Vitavax® 25-50 W.P. Seed Protectant: S-806: SOCAL 711/XXII:14. (Unpublished study received Jul 22, 1975 under 239-2451; prepared by Standard Oil Co. of California, Safety Div., submitted by Chevron Chemical Co., Richmond, Calif.; CDL:224836-D)

NRID	CITATION
00002887	Rittenhouse, J.R.; Narcisse, J.K. (1975) The Skin Irritation Potential of Difolatan-Vitavax® 1-2F Seed Protectant (CC 5640) and Difolatan-Vitavax® 1-2F Seed Protectant (CC 5998): S-801: SOCAL 705/XXII:17. (Unpublished study received Jul 22, 1975 under 239-2448; prepared by Standard Oil Co. of California, Safety Div., submitted by Chevron Chemical Co., Richmond, Calif.; CDL: 230799-E)
00002891	Rittenhouse, J.R.; Narcisse, J.K. (1975) The Skin Irritation Potential of Orthocide-Vitavax® 25-50 W.P. Seed Protectant: S-807: SOCAL 712/XXII:15. (Unpublished study received Jul 22, 1975 under 239-2451; prepared by Standard Oil Co. of California, Safety Div., submitted by Chevron Chemical Co., Richmond, Calif.; CDL:224836-E)
00003027	Rittenhouse, J.R.; Narcisse, J.K. (1975) The Skin Sensitization Potential of Orthocide-Vitavax® 25-50 W.P. Seed Protectant (CC 6033): S-809: SOCAL 714/XXIV:41. Rev. (Unpublished study received May 11, 1977 under 239-2458; prepared by Standard Oil Co. of California, submitted by Chevron Chemical Co., Richmond, Calif.; CDL:230224-G)
00002878	Rittenhouse, J.R.; Narcisse, J.K. (1976) The Skin Sensitization Potential of Difolatan-Vitavax® 1-2F Seed Protectant (CC 5998): S-803: SOCAL 707B/XXIV:38. Rev. (Unpublished study received Jul 6, 1977 under 239-2459; prepared by Standard Oil Co. of California, Safety and Health Div., submitted by Chevron Chemical Co., Richmond, Calif.; CDL:230827-G)
00025475	Roberts, T.G. (1969) [Efficacy Data for Vitavax® on Peanuts]. (Unpublished study received Jan 12, 1970 under 0F0939; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091604-N)
00003038	Robinson, D.; Shillam, K.W.G. (1968) The Palatability to the Young Pheasant of Grain Dressed with Vitavax® and Its Effects on Mortality and Growth: 2469/68/346. (Unpublished study received Jun 14, 1969 under 9G0819; prepared by Huntingdon Research Centre, Eng. for Uniroyal International, submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091420-P)
05005920	Schewe, T.; Hiebsch, C. (1977) Zur Wirkung von Atmungshemmstoffen auf das Elektronentransportsystem von <i>Escherichia coli</i> . [Effect of respiratory inhibitors on the electron transport system of <i>Escherichia coli</i> .] <i>Acta Biologica et Medica Germanica</i> 36(7/8):961-966.
05013093	Schewe, T.; Hiebsch, C.; Garcia Parra, M.; Rapoport, S. (1974) Zur Wirkung von Atmungshemmstoffen auf die Atmungsenzyme der Mitochondrien aus Blumenkohl (<i>Brassica oleracea</i> L.). [The effect of respiratory inhibitors upon the respiratory enzymes of mitochondria from cauliflower (<i>Brassica oleracea</i> L.).] <i>Acta Biologica Et Medica Germanica</i> 32(5):419-426.

MRID

CITATION

- 05013096 Schewe, T.; Rapoport, S.; Boehme, G.; Kunz, W. (1975) Zur Wirkung von Carboxin auf das Elektronentransportsystem tierischer Mitochondrien. [The effects of carboxin on the electron transport system of animal mitochondria.] Pages 145-152, In Systemfungizide, Internationales Symposium; May, 1974, Reinhardtsbrunn, East Germany. Edited by H. Lyr and C. Polter. Berlin, East Germany: Akademie-Verlag.
- 05015868 Schewe, T.; Rapoport, S.; Boehme, J.; Kunz, W. (1973) Zum Angriffspunkt des Systemfungizids Carboxin in der Atmungskette. [The site of action of the systemic fungicide carboxin in the respiratory chain.] Acta Biologica Et Medica Germanica 31(1):73-86.
- 00003117 Shapiro, R. (1977) [Toxicity Data--Vitavax®]: Report No. T-272. (Unpublished study received Jan 24, 1978 under 400-81; prepared by Product Safety Labs, Div. of Nutrition International, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:233500-C)
- 05002742 Sikka, V.K.; Chopra, B.; Sethi, R.K.; Chopra, S.L. (1973) Persistence of Vitavax® on (5,6-dihydro-3-carboxanilidro-2 methyl-1, 4-oxathiin) residues on some crops. Pesticides 7(4):28-29.
- 05009406 Sikka, V.K.; Chopra, B.; Sethi, R.K.; Chopra, S.L. (1973) Persistence of Vitavax® on (5,6-dihydro-3-carboxanilidro-2 methyl-1,4-oxathiin) residues on some crops. Pesticides VII(4):28-29.
- 05003526 Sinha, O.K.; Vyas, S.C.; Joshi, L.K. (1975) Rapid separation and detection of some systemic fungicides by paper chromatography. Journal of Chromatography 110(2):403-405.
- 05004768 Sinha, S.K.; Prasad, M. (1977) Studies on certain aspects of chemical control of bacterial stalk rot disease of maize. Zentralblatt fuer Bakteriologie, Parasitenkunde, Infektionskrankheiten und Hygiene, Abteilung 2 132(1):89-92.
- 00002938 Sisken, H.R. (1970) Determination of Residual Vitavax® in Meal and Oil from Cotton and Peanut Seed. (Unpublished study received Jul 28, 1972 under OF0939; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091603-A)
- 00025465 Sisken, H.R. (1970) Determination of Vitavax® and its Sulfoxide in Cotton Seed: Modification I of General GLC Method. Method dated Jan 5, 1970. (Unpublished study received Jan 12, 1970 under OF0939; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091604-C)
- 00025466 Sisken, H.R.; Lane, J.R. (1970) Storage Stability Studies of Vitavax® Residue in Grain. (Unpublished study received Jan 12, 1970 under OF0939; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091604-E)

MRID	CITATION
00003184	Sisken, H.R.; Newell, J.E. (1969) The Determination of Residual Vitavax [®] and F-831 in Wheat Seeds: Tentative Method. Method dated Feb 20, 1969. (Unpublished study received Mar 25, 1969 under 9G0819; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:093520-AB)
00003052	Sisken, H.R.; Newell, J.E. (1970) General GLC Method for Determination of Residual Vitavax [®] and Its Sulfoxide in Seeds (Grain and Peanuts). Method dated Jan 5, 1970. (Unpublished study received Jan 12, 1970 under 0F0939; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:093245-Q)
00003335	Sisken, H.R.; Newell, J.E. (1971) Determination of residues of Vitavax [®] and its sulfoxide in seeds. Journal of Agricultural and Food Chemistry 19(4):738-741. (Also?In?unpublished submission received Apr 21, 1977 under unknown admin. no.; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:229370-P)
00003088	Smilo, A.R.; Lacadie, J.A.; Cardona, R. (1977) Photochemical Fate of Vitavax [®] in Solution. (Unpublished study received May 3, 1977 under 400-80; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:231932-C)
05003663	Snel, M.; Edgington, L.V. (1970) Uptake, translocation and decomposition of systemic oxathiin fungicides in bean. Phytopathology 60(12):1708-1716.
05003661	Solel, Z.; Edgington, L.V. (1973) Transcuticular movement of fungicides. Phytopathology 63(4):505-510.
0500444	Solel, Z.; Pinkas, J. (1972) Bioassay evaluation of the systemic properties of carboxin in lemon seedlings. Pesticide Science 3(2):149-151.
05003657	Staphorst, J.L.; Strijdom, B.W. (1976) Effects on rhizobia of fungicides applied to legume seed. Phytophylactica 8(2):47-54.
00005856	Stevens, K.R. (1979) Primary Skin Irritation Study with Rabbits. (Unpublished study received Feb 28, 1979 under 400-124; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:237477-B)
00024924	Stevens, K.R. (1979) Report: [Vitavax 30-C]. (Unpublished study received Aug 24, 1979 under 400-124; prepared by Food and Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:240917-C)
00005858	Stevens, K.R. (1979) Report: Acute Dermal Toxicity Study in Rabbits. (Unpublished study received Feb 28, 1979 under 400-124; prepared by Food & Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:237477-D)

MRID	CITATION
00005857	Stevens, K.R. (1979) Report: Eye Irritation Test in Rabbits. (Unpublished study received Feb 28, 1979 under 400-124; prepared by Food & Drug Research Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:237477-C)
05005314	Stoker, G.L.; Dewey, W.G. (1969) Effects of loose smut infection and Vitavax [®] seed treatment on barley yields. Crop Science 9(5):586-588.
00025472	Stoker, G.L.; Dewey, W.G. (1969) Effects of loose smut infection and Vitavax [®] seed treatment on barley yields. Crop Science 9 (Sep-Oct):586-588. (Also In unpublished submission received Jan 12, 1970 under 0F0939; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091604-K)
GS001201	Stone, G.M. (1969) Stability of Vitavax [®] in Stored Plants. (Unpublished study received Jan 12, 1970 under 0F0939; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091604)
00003048	Stone, G.M. (1971) Concentration of Vitavax [®] (PPM) on Treated Seed. (Unpublished study received on unknown date under 0F0939; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:093245-II)
00003050	Stone, G.M. (1971) Quantity of Vitavax [®] in Soil around the Seed. (Unpublished study received on unknown date under 0F0939; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:093245-J)
05003853	Stone, G.M. (1976) Vitavax [®] . Pages 319-331, In Analytical Methods for Pesticides and Plant Growth Regulators. Vol. 8. New York: Academic Press.
00025464	Stone, G.M.; Chin, W.T.; Smith, A.E. (1970) Residues of Vitavax [®] (Carboxin) in Plants: 14C Studies. (Unpublished study received Jan 12, 1970 under 0F0939; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091604-B)
05003969	Swinburne, T.R. (1975) The effect of benomyl and other fungicides on phenylalanine-ammonia lyase activity in <u>Hordeum vulgare</u> and <u>Phaseolus vulgaris</u> . Physiological Plant Pathology 5(1):81-88.
05002994	Thorn, G.D.; Richardson, L.T. (1970) Effects of various fungicides in soil on water and amino acid transport in tomato plants. Canadian Journal of Botany 48(11):2033-2036.
05018561	Thornburg, W. (1971) Pesticide residues. Analytical Chemistry 43(5):145-162.
05002150	Thornburg, W. (1973) Pesticide residues. Analytical Chemistry 45(5):151R-167R.
05009700	Thornburg, W. (1973) Pesticide residues. Analytical Chemistry 45(5):151-167.

MRID	CITATION
05003435	Tripathi, R.K.; Bhaktavatsalam, G. (1973) Silver nitrate-bromophenol blue and potassium permanganate-sulphuric acid as chromogenic reagents for the detection of carboxin and oxycarboxin in thin-layer chromatography. <i>Journal of Chromatography</i> 87(1):283-286.
05004133	Tripathi, R.K.; Singh, A.; Singh, S.L. (1976) Residue analysis of wheat grains and straw produced from Vitavax-treated seed. <i>Pesticides</i> 10(8):33-34.
05002996	Tucker, A.N.; Lillich, T.T. (1974) Effect of the systemic fungicide carboxin on electron transport function in membranes of <u>Micrococcus denitrificans</u> . <i>Antimicrobial Agents and Chemotherapy</i> 6(5):572-578.
00003138	Uniroyal (19??) D735 and F461 Systemic Fungicides: Technical Data Sheet. (Unpublished study received on unknown date under unknown admin. no.; CDL:125233-A)
00003039	Uniroyal Chemical (19??) [Residue Studies: Vitavax]. (Unpublished study including summary, received Jun 14, 1969 under 9G0819; CDL:091420-R)
00003160	Uniroyal Chemical (19??) [Summaries for "Handling" and "Fish and Wildlife Toxicology"]. Summary of studies 229370-C through 229370-N. (Unpublished study received Apr 21, 1977 under unknown admin. no.; CDL:229370-B)
00002920	Uniroyal Chemical (19??) Calculation of Margin of Safety: [Vitavax®]. (Unpublished study received Jun 9, 1975 under 5F1637; CDL:094947-L)
00002860	Uniroyal Chemical (19??) Chemical and Physical Properties of Vitavax® Technical: Technical Bulletin. (Unpublished study received Sep 17, 1976 under 400-132; CDL:226141-B)
00021623	Uniroyal Chemical (19??) Chemical and Physical Properties: Vitavax®. (Unpublished study received Jul 19, 1968 under 400-80; CDL:003258-A)
00025479	Uniroyal Chemical (19??) Chemical and Physical Properties: Vitavax®. (Unpublished study received Jun 17, 1974 under 5F1525; CDL:094043-A)
00003171	Uniroyal Chemical (19??) Chemical and Physical Properties: vitavax® (Unpublished study received Mar 25, 1969 under 9G0819; CDL:093520-D)
00002995	Uniroyal Chemical (19??) Control Test AC-1045-D: Assay of Vitavax® (and Its Formulations). Undated method. (Unpublished study received Jul 15, 1976 under 400-118; CDL:224929-C)

MRID	CITATION
00024926	Uniroyal Chemical (19??) Control Test: AC-1045-D: Assay of Vitavax®. (and Its Formulations). Undated method. (Unpublished study received Mar 27, 1975 under 0F0939; CDL:098575-A)
00003086	Uniroyal Chemical (19??) Environmental Fate Studies on Vitavax®. (Unpublished study received May 3, 1977 under 400-80; CDL:231932-A)
00004159	Uniroyal Chemical (19??) Margin of Safety of Proposed Tolerances: [Vitavax®]. (Unpublished study received Jan 12, 1970 under 0F0939; CDL:093245-R)
00004158	Uniroyal Chemical (19??) Margin of Safety of the Proposed Tolerances: [Vitavax®]. (Unpublished study received May 1, 1974 under 4F1499; CDL:094551-I)
00030598	Uniroyal Chemical (19??) Margin of Safety of the Proposed Tolerances: [Vitavax®]. (Unpublished study received Dec 23, 1972 under 3F1318; CDL:092253-A)
00003232	Uniroyal Chemical (19??) Residue Summary for Carboxin in Peanuts. (Unpublished study received Mar 20, 1979 under 400-80; CDL:098014-C)
00002948	Uniroyal Chemical (19??) Seed Rots and Seedling Disease Control Data Summary. (Unpublished study received Dec 23, 1972 under 3F1318; CDL:092253-B)
00003103	Uniroyal Chemical (19??) Stability of Vitavax® on Treated Oat Seed. (Unpublished study received Apr 21, 1975 under 400-112; CDL:220785-A)
00003092	Uniroyal Chemical (19??) Stability of Vitavax® on Treated Wheat and Barley Seed. (Unpublished study received Jun 26, 1973 under 400-108; CDL:008327-D)
00002924	Uniroyal Chemical (19??) Summary of Active Handling Toxicology--Use Category, Signal Word and Precautionary Statements. (Unpublished study received Sep 28, 1976 under 400-118; CDL:230403-A)
00025485	Uniroyal Chemical (19??) Summary of Field Residue Data. (Unpublished study received Jan 12, 1970 under 0F0939; CDL:091604-R)
00003123	Uniroyal Chemical (19??) Summary of Phytotoxicity Data. (Unpublished study received May 16, 1973 under 400-107; CDL:003284-D)
00002959	Uniroyal Chemical (19??) Summary: [Vitavax® Residue]. (Unpublished study received Jul 28, 1972 under 0F0939; CDL:091603-B)
00002971	Uniroyal Chemical (19??) The Results of Tests on the Amount of Residue Remaining, Including a Description of the Analytical Method Used. (Unpublished study received on unknown date under 9G0819; CDL:091418-L)

MRID	CITATION
00021616	Uniroyal Chemical (19??) Vitavax® HBM-25 Production Procedure. (Unpublished study received Apr 25, 1979 under 400-136; CDL: 238225-A)
00029651	Uniroyal Chemical (19??) Vitavax® HBM-25 Production Procedure. (Unpublished study received Apr 25, 1979 under 400-136; CDL: 238225-A)
00003170	Uniroyal Chemical (19??) Vitavax® Manufacturing Process. (Unpublished study received Mar 25, 1969 under 9G0819; CDL:093520-C)
00021644	Uniroyal Chemical (19??) Vitavax® Manufacturing Process. (Unpublished study received Jul 19, 1968 under 400-80; CDL:003258-AN)
00003296	Uniroyal Chemical (19??) Vitavax®: Chemical and Physical Properties of Vitavax® Technical. (Unpublished study received Sep 27, 1972 under 3F1318; CDL:093547-J)
00003231	Uniroyal Chemical (19??) Vitavax® Flowable Fungicide: Vitavax®-3F Fungicide: Basic Manufacturing Process. (Unpublished study received Mar 20, 1979 under 400-80; CDL:098014-B)
00003230	Uniroyal Chemical (19??) Vitavax® Fungicide: Agricultural Chemicals: Technical Data Sheet. (Unpublished study received Mar 20, 1979 under 400-80; CDL:098014-A)
00005860	Uniroyal Chemical (19??) Vitavax®-4G Process. (Unpublished study received Feb 6, 1979 under 400-EX-58; CDL:237329-B)
00002979	Uniroyal Chemical (1962?) Margin of Safety of the Proposed Tolerances: [Vitavax®]. (Unpublished study received Aug 11, 1971 under 2F1191; CDL:093516-H)
00003055	Uniroyal Chemical (1962) Calculation of Margin of Safety: [Vitavax®] (Unpublished study received Jun 9, 1975 under 5F1638; CDL:094948-H)
00002978	Uniroyal Chemical (1966) Vitavax® Assay Method. Method dated Jan 24, 1966. (Unpublished study received Aug 11, 1971 under 2F1191; CDL:093516-F)
00003185	Uniroyal Chemical (1967) Residue Analysis of Cotton Seedlings from Vitavax®-Treated Cotton Seeds. (Unpublished study received Nov 29, 1967 under 400-EX-28; CDL:123430-A)
00021632	Uniroyal Chemical (1968) Reasonable Grounds for Support of This Petition: [Vitavax®]. (Unpublished study received Jul 19, 1968 under 400-80; CDL:003258-R)
00021629	Uniroyal Chemical (1968) The Results of Tests on the Amount of Residue Remaining, Including a Description of the Analytical Method Used: [Vitavax®]. (Unpublished study received Jul 19, 1968 under 400-80; CDL:003258-N)

MRID	CITATION
00005538	Uniroyal Chemical (1968) Vitavax®-Fungicide: Peanut Seed Treatment Test Summary in Combination with Difolatan. (Unpublished study received May 2, 1975 under 400-81; CDL:098028-I)
00003131	Uniroyal Chemical (1969?) Basic Vitavax® Toxicology Studies. (Unpublished study received May 16, 1973 under 400-107; CDL: 003284-R)
00003072	Uniroyal Chemical (1969?) Report of Investigations Made with Respect to the Safety of Vitavax®. (Unpublished study received Mar 25, 1969 under 400-EX-38; CDL:127028-A)
00002946	Uniroyal Chemical (1969?) Uniroyal Supplemental Experimental Studies: Animal Residues. (Unpublished study received Oct 9, 1973 under 3F1318; CDL:092254-K)
00003051	Uniroyal Chemical (1969) Data on Seed Treatment--Soil Fungicide Test for Control of Cotton Seedling Diseases, Athens, Georgia, 1969: Table 2. (Unpublished study received Jun 5, 1970 under OF0939; CDL:093245-N)
00003045	Uniroyal Chemical (1969) Disappearance Studies: Wheat and Barley; [Peanut; Sorghum: Vitavax®]. (Unpublished study received Jun 14, 1969 under 9G0819; CDL:091420-X)
00003029	Uniroyal Chemical (1969) Toxicology Information: [Vitavax®]. (Unpublished study received Jun 14, 1969 under 9G0819; CDL: 091420-A)
00005539	Uniroyal Chemical (1969) Vitavax®-Fungicide: Peanut Seed Treatment Test Summary in Combination with Botran. (Unpublished study received May 2, 1975 under 400-81; prepared in cooperation with Oklahoma State Univ. and North Carolina State Univ.; CDL: 098028-J)
00003307	Uniroyal Chemical (1970?) Additional Vitavax® Toxicology. Summary of studies 091003-C through 091003-R. (Unpublished study received May 8, 1972 under 2F1191; CDL:091003-E)
00003057	Uniroyal Chemical (1970?) Appendix. (Unpublished study received Dec 21, 1971 under 400-81; CDL:050050-A)
00025481	Uniroyal Chemical (1970) Basic Vitavax® Toxicology Studies. (Unpublished study received Jun 17, 1974 under 5F1525; CDL:094043-E)
00002939	Uniroyal Chemical (1970) Vitavax® Handling Toxicity Studies. (Unpublished study received Oct 9, 1973 under 3F1318; CDL:092254-C)
00003132	Uniroyal Chemical (1971?) Toxicity of Vitavax® to Fish and Wildlife. (Unpublished study received May 16, 1973 under 400-107; CDL: 003284-S)

MRID	CITATION
00003073	Uniroyal Chemical (1971?) Toxicity to Fish and Wildlife: [Vitavax®]. (Unpublished study received on unknown date under 400-EX-38; CDL:127028-B)
00002962	Uniroyal Chemical (1971) [Residue Data]. (Unpublished study received May 8, 1972 under 2F1191; CDL:091003-S)
00003078	Uniroyal Chemical (1971) Background of Residue Data for Vitavax®-200 Fungicide and Vitavax®-300 Fungicide on Peanuts. (Unpublished study received Jun 10, 1975 under 400-92; CDL:220663-J)
00002982	Uniroyal Chemical (1971) Composition of Vitavax®--Technical Grade. (Unpublished study received on July 10, 1974 under 2F1525; CDL:095561-B)
00002989	Uniroyal Chemical (1971) Cotton Seedling Survival. (Unpublished study received Apr 12, 1972 under 400-80; CDL:023352-D)
00002917	Uniroyal Chemical (1971) Peanuts: Seed Treatment--1971. (Unpublished study received May 1, 1974 under 4F1499; CDL:094551-X)
00003089	Uniroyal Chemical (1971) Peanuts: Seed Treatment--1971: Jackson, N.C. (Unpublished study received Jun 10, 1975 under 400-92; CDL:220663-O)
00002859	Uniroyal Chemical (1971) Reasoning in Support of the Petition: [Vitavax®]. (Unpublished study including revised section G-5, received Sep 27, 1972 under 3F1318; CDL:093547-O)
00002862	Uniroyal Chemical (1971) Regional Pod Rot Test--1971. (Unpublished study received Sep 28, 1976 under 400-130; CDL:230405-O)
00002859	Uniroyal Chemical (1971) Reasoning in Support of the Petition: received Sep 27, 1972 under 3F1318; CDL:093547-O)
00002954	Uniroyal Chemical (1971) Seed Corn Treatment Experiment, Washington, Iowa. (Unpublished study received Dec 23, 1972 under 3F1318; CDL:092253-K)
00005536	Uniroyal Chemical (1971) Vitavax®--Fungicide: Peanut Seed Treatment Test Summary in Combination with Captan. (Unpublished study received May 2, 1975 under 400-81; prepared in cooperation with Auburn Univ., Oklahoma State Univ. and Texas A & M Univ., Agricultural Extension Service; CDL:098028-G)
00005537	Uniroyal Chemical (1971) Vitavax®--Fungicide: Peanut Seed Treatment Test Summary in Combination with Thiram. (Unpublished study received May 2, 1975 under 400-81; CDL:098028-H)
00002942	Uniroyal Chemical (1972) [Animal Residue Studies]. (Unpublished study received Oct 9, 1973 under 3F1318; CDL:092254-G)

MRID	CITATION
00002961	Uniroyal Chemical (1972) [Residue Data: Vitavax®]. (Unpublished study received on unknown date under 2F1191; prepared in cooperation with Morse Laboratories and Harris Laboratories; CDL:091003-A)
00002951	Uniroyal Chemical (1972) Field Performance Report. (Unpublished study including report, received Dec 23, 1972 under 3F1318; CDL:092253-F)
00003125	Uniroyal Chemical (1972) Field Performance Report. (Unpublished study received May 16, 1973 under 400-107; CDL:003284-H)
00002951	Uniroyal Chemical (1972) Field Performance Report. (Unpublished study including report, received Dec 23, 1972 under 3F1318; CDL:092253-F)
00003128	Uniroyal Chemical (1972) Field Phytotoxicity Evaluation. (Unpublished study received May 16, 1973 under 400-107; CDL:003284-N)
00002864	Uniroyal Chemical (1972) Pod Rot Test--1972. (Unpublished study received Sep 28, 1976 under 400-130; CDL:230405-R)
00002928	Uniroyal Chemical (1972) Residues in PPM: [Vitavax®]. (Unpublished study received on unknown date under 0F0939; CDL:094582-B)
00002930	Uniroyal Chemical (1972) Residues in PPM: [Vitavax®]. (Unpublished study received on unknown date under 0F0939; CDL:094582-E)
00002858	Uniroyal Chemical (1972) Stability of Vitavax® on Treated Seed (Grain and Corn). (Unpublished study received Sep 27, 1972 under 3F1318; CDL:093547-M)
00003304	Uniroyal Chemical (1972) Stability of Vitavax®-200 in Treated Seed (Cotton). (Unpublished study received Jun 28, 1972 under 0F0939; CDL:094582-F)
00003303	Uniroyal Chemical (1972) Stability of Vitavax®-300 in Treated Seed (Cotton). (Unpublished study received Jun 28, 1972 under 0F0939; CDL:094582-D)
00002940	Uniroyal Chemical (1972) Summary of Vitavax® Residue Data in Corn and Small Grains. Includes method dated Nov 1, 1968. (Unpublished study including report, received Oct 9, 1973 under 3F1318; CDL:092254-E)
00002902	Uniroyal Chemical (1972) Untreated Peanuts Analyzed by Vitavax® (GLC) Method for Apparent PPM Background. (Unpublished study received on unknown date under 4F1499; CDL:093979-B)
00025486	Uniroyal Chemical (1972) Vitavax® Handling Toxicity Studies. (Unpublished study received Jun 17, 1974 under 5F1525; CDL:094043-J)

MRID	CITATION
00002960	Uniroyal Chemical (1973) Discussion: [Vitavax® Residues]. (Unpublished study received Oct 9, 1973 under 3F1318; CDL:092254-M)
00002857	Uniroyal Chemical (1973) Enforcement Method for Vitavax® in Animal Tissue, Milk and Eggs. Method dated Sep 17, 1973. (Unpublished study received on unknown date under 3F1318; CDL:093547-D)
00003155	Uniroyal Chemical (1973) Field Performance and Phytotoxicity Report. (Unpublished study received Dec 5, 1973 under 400-107; CDL:023365-H)
00003094	Uniroyal Chemical (1973) Field Performance Report: UNI-1088 and 1090. (Unpublished study received Jun 26, 1973 under 400-108; CDL:008327-F)
00002910	Uniroyal Chemical (1973) Field Performance and Phytotoxicity Report. (Unpublished study received May 1, 1974 under 4F1499; CDL:094551-P)
00003305	Uniroyal Chemical (1973) Method for Vitavax® Residues in Animal Tissue, Milk and Eggs. Method dated Sep 17, 1973. (Unpublished study received Oct 9, 1973 under 3F1318; CDL:092254-A)
00002908	Uniroyal Chemical (1973) Peanut Seed Treatment 1973: 50 Seed Planted/Plot. (Unpublished study received May 1, 1974 under 4F1499; CDL:094551-M)
00003129	Uniroyal Chemical (1973) Residues in PPM. (Unpublished study received May 16, 1973 under 400-107; prepared in cooperation with Morse Laboratories, Inc.; CDL:003284-P)
00005852	Uniroyal Chemical (1973) Residues in PPM: Corn: UNI-1088. (Unpublished study received Mar 28, 1977 under 400-107; prepared in cooperation with New York State Univ.--Oswego, Lake Ontario Environmental Laboratory and Morse Laboratories, Inc.; CDL: 238083-A)
00005854	Uniroyal Chemical (1973) Residues in PPM: Wheat & Barley. (Unpublished study received Feb 28, 1978 under 400-115; CDL:238076-E)
00003121	Uniroyal Chemical (1973) Stability of Vitavax® on Treated Cotton Seed. (Unpublished study received May 16, 1973 under 400-107; CDL:003284-B)
00003158	Uniroyal Chemical (1973) Vitavax® Flowable Fungicide: Summary of Residues in Wheat and Barley. (Unpublished study received Dec 5, 1973 under 400-107; CDL:023365-S)
00005530	Uniroyal Chemical (1973) Vitavax®--Fungicide: Peanut Seed Treatment Test Summary at 3-6 Oz/100 Pounds of Seed. (Unpublished study received May 2, 1975 under 400-81; CDL:098028-A)

MRID	CITATION
00003159	Uniroyal Chemical (1973) Vitavax [®] -200 Flowable Fungicide: Summary of Residues in Wheat and Barley. (Unpublished study received Dec 5, 1973 under 400-107; CDL:023365-T)
00002861	Uniroyal Chemical (1973) Vitavax [®] 10G: Peanuts: Phytotoxicity. (Unpublished study received Sep 28, 1976 under 400-130; CDL: 230405-N)
00003106	Uniroyal Chemical (1974) Field Evaluation Report: Phyto Study. (Unpublished study received Apr 21, 1975 under 400-112; CDL: 220785-D)
00003001	Uniroyal Chemical (1974) Phytotoxicity. (Unpublished study received Jul 15, 1976 under 400-118; CDL:224933-B)
00002903	Uniroyal Chemical (1974) Residue Analysis of Peanut Hulls for Vitavax [®] . (Unpublished study received on unknown date under 4F1499; CDL:093979-C)
00002904	Uniroyal Chemical (1974) Residue Analysis of Peanut Meal for Vitavax [®] . (Unpublished study received on unknown date under 4F1499; CDL:093979-D)
00002937	Uniroyal Chemical (1974) Residues in PPM: [Vitavax [®]]. (Unpublished study received Mar 21, 1974 under 400-113; prepared in cooperation with Morse Laboratories, Inc.; CDL:023366-C)
00003021	Uniroyal Chemical (1974) Residues in PPM: Soybeans: Vitavax [®] . (Unpublished study received Nov 14, 1978 under 400-112; CDL:235936-W)
00005848	Uniroyal Chemical (1974) Residues in PPM: Soybeans: Vitavax [®] UNI-1127. (Unpublished study received Dec 11, 1977 under 400-112; CDL:238081-E)
00002905	Uniroyal Chemical (1974) The Effects of EPA Tolerance Pesticides upon the Recovery of Vitavax [®] Residues from Peanuts. (Unpublished study received on unknown date under 4F1499; prepared in cooperation with Morse Laboratories, Inc.; CDL:093979-F)
00002901	Uniroyal Chemical (1974) The Results of Tests on the Amount of Residue Remaining, Including a Description of the Analytical Methods Used: [Vitavax [®]]. Summary of studies 093979-B through 093979-F. (Unpublished study including summary, received on unknown date under 4F1499; CDL:093979-A)
00025482	Uniroyal Chemical (1974) Toxicity of Vitavax [®] to Fish and Wildlife. (Unpublished study received Jun 17, 1974 under 5F1525; CDL: 094043-G)
00002957	Uniroyal Chemical (1974) Toxicity of Vitavax [®] to Fish and Wildlife. (Unpublished study received Mar 21, 1974 under 400-113; CDL: 023366-F)

MRID	CITATION
00003002	Uniroyal Chemical (1975?) Summary of Acute Handling Toxicology--Use Category, Signal Word and Precautionary Statements. Summary of study 224934-C. (Unpublished study received Jul 15, 1976 under 400-118; CDL:224934-B)
00003003	Uniroyal Chemical (1975?) Summary of Fish and Wildlife Toxicology for Precautionary Label Statements. Summary of study 224935-C. (Unpublished study received Jul 15, 1976 under 400-118; CDL: 224935-B)
00003004	Uniroyal Chemical (1975?) Toxicity of Vitavax [®] to Fish and Wildlife. (Unpublished study received Jul 15, 1976 under 400-118; CDL:224935-C)
00002871	Uniroyal Chemical (1975?) Toxicity of Vitavax [®] to Fish and Wildlife. (Unpublished study received Sep 28, 1976 under 400-130; CDL: 230406-C)
00024930	Uniroyal Chemical (1975) [Residues from Vitavax [®] on Cotton]. (Unpublished study received Mar 27, 1975 under 0F0939; prepared in cooperation with Morse Laboratories, Inc.; CDL:098575-F)
00003149	Uniroyal Chemical (1975) Field Evaluation Report. (Unpublished study received Sep 13, 1976 under 400-129; CDL:225604-W)
00003356	Uniroyal Chemical (1975) Residues of Vitavax [®] --EVS Concentrate in Corn. (Unpublished study received Jun 10, 1975 under 400-124; prepared in cooperation with Morse Laboratories, Inc.; CDL:235928-C)
00021647	Uniroyal Chemical (1975) Response to Question No. 5 Concerning Status Report No. 1,...on the Environmental Fate of Vitavax [®] . (Unpublished study received on unknown date under 400-80; CDL:240822-A)
00002918	Uniroyal Chemical (1975) Stability of Vitavax [®] on Treated Soybean Seed. (Unpublished study received Jun 9, 1975 under 5F1637; CDL:094947-E)
00003108	Uniroyal Chemical (1975) Summary of Residue Data: [Vitavax [®]]. (Unpublished study received Apr 21, 1975 under 400-112; prepared in cooperation with Morse Laboratories, Inc.; CDL:220785-G)
00003054	Uniroyal Chemical (1975) The Effects of EPA Tolerance Pesticides upon the Recovery of Vitavax [®] Residues from Sorghum: (1) Grain, (2) Fodder and Forage. (Unpublished study including summary, received Jun 9, 1975 under 5F1638; prepared in cooperation with Morse Laboratories, Inc.; CDL:094948-G)
00002919	Uniroyal Chemical (1975) The Effects of EPA Tolerance Pesticides upon the Recovery of Vitavax [®] Residues from Soybeans. (Unpublished study including summary, received Jun 9, 1975 under 5F1637; prepared in cooperation with Morse Laboratories, Inc.; CDL:094947-K)

MRID	CITATION
00003017	Uniroyal Chemical (1976?) Soybeans: Vitavax®-200 Flowable Fungicide. (Unpublished study received Nov 14, 1978 under 400-12; CDL:235936-D)
00003130	Uniroyal Chemical (1976?) Vitavax® Handling Toxicity Studies. (Unpublished study including annotated bibliography of petition materials, received May 16, 1973 under 400-107; CDL:003284-Q)
00003085	Uniroyal Chemical (1976) [Vitavax® Residue Data on Peanuts]. (Unpublished study received Sep 13, 1976 under 400-129; CDL:225602-A)
00003084	Uniroyal Chemical (1976) Vitavax® Manufacturing Process. (Unpublished study including letter dated Sep 9, 1976 from T.E. Geise to John Pryzbylek, received Sep 13, 1976 under 400-129; CDL:225601-A)
00005862	Uniroyal Chemical (1977?) Toxicity: Vitavax®. (Unpublished study received Feb 6, 1979 under 400-EX-58; CDL:237329-D)
00003115	Uniroyal Chemical (1977?) Toxicology: Vitavax® Technical. Summary of studies 233500-B through 233500-F. (Unpublished study received Jan 24, 1978 under 400-81; CDL:233500-A)
00003009	Uniroyal Chemical (1977?) Toxicology: Vitavax®-200 Flowable. (Unpublished study received Jul 14, 1978 under 400-112; CDL:235042-A)
00003165	Uniroyal Chemical (1977) [Residue Data: Vitavax®]. (Unpublished study received Apr 21, 1977 under unknown admin. no.; prepared in cooperation with Morse Laboratories, Inc.; CDL:229370-O)
00003357	Uniroyal Chemical (1977) Chemical and Physical Properties of Vitavax® Technical. (Unpublished study received Oct 11, 1978 under KS-78/20; submitted by ?; CDL:235366-A)
00003234	Uniroyal Chemical (1977) Reasonable Grounds in Support of This Petition: [Vitavax®]. (Unpublished study received Mar 20, 1979 under 400-80; CDL:098014-F)
00020892	Uniroyal Chemical (1977) Residues in PPM. (Unpublished study received Apr 25, 1979 under ND 79/8; CDL:238250-E)
00003221	Uniroyal Chemical (1977) Residues in PPM: Barley: Vitavax® 25DB: Grain & Straw. (Unpublished study received May 10, 1978 under 400-115; prepared in cooperation with Morse Laboratories, Inc.; CDL:235653-E)
00030107	Uniroyal Chemical (1977) Toxicology Vitavax HBM-25. Summary of studies 238226-B through 238286-F. (Unpublished study received Apr 25, 1979 under 400-136; CDL:238226-A)

MRID	CITATION
00003079	Uniroyal Chemical (1977) Toxicology: Vitavax® Flowable Fungicide. (Unpublished study received Dec 29, 1977 under 400-107; CDL: 232545-A)
00005863	Uniroyal Chemical (1977) Toxicology: Vitavax® 10G. Summary of studies 237329-F through 237329-I. (Unpublished study received Feb 6, 1979 under 400-EX-58; CDL:237329-E)
00005859	Uniroyal Chemical (1977) Vitavax® Fungicide: Technical Data Sheet. (Unpublished study received Feb 6, 1979 under 400-EX-58; CDL:237329-A)
00020891	Uniroyal Chemical (1977) Vitavax®-25 DP: Phytotoxicity. (Unpublished study received Apr 25, 1979 under ND 79/8; prepared in cooperation with Univ. of Idaho, Experiment Station; CDL: 238250-D)
00003236	Uniroyal Chemical (1977) Vitavax®-3F: Peanuts. (Unpublished study received May 6, 1978 under 400-EX-55; CDL:234122-B)
00005548	Uniroyal Chemical (1977) Vitavax®-25DB: Phytotoxicity. (Unpublished study received Feb 28, 1978 under 400-115; prepared in cooperation with the University of Idaho, Experiment Station; CDL:238076-D)
00024920	Uniroyal Chemical (1978) [Vitavax® Residues in Wheat, Barley and Oat]. (Unpublished study received Sep 26, 1979 under 400-107; prepared in cooperation with Morse Laboratories, Inc.; CDL: 241014-A)
00003220	Uniroyal Chemical (1978) Residues in PPM: Oats: Vitavax® 25DB: Grain & Straw. (Unpublished study received May 10, 1978 under 400-115; prepared in cooperation with Morse Laboratories, Inc.; CDL:235653-D)
00003219	Uniroyal Chemical (1978) Residues in PPM: Wheat: Vitavax® 25DB: Forage. (Unpublished study received May 10, 1978 under 400-115; prepared in cooperation with Morse Laboratories, Inc.; CDL: 235653-C)
00003218	Uniroyal Chemical (1978) Residues in PPM: Wheat: Vitavax® 25DB: Grain & Straw. (Unpublished study received May 10, 1978 under 400-115; prepared in cooperation with Morse Laboratories, Inc.; CDL:235653-B)
00003224	Uniroyal Chemical (1978) Response to EPA Letter Dated 6 October, 1977: Environmental Fate of Vitavax® Fungicide. Summary of studies 236662-F through 236662-J. Includes method no. 2344 dated Aug 22, 1978. (Unpublished study received Dec 27, 1978 under 400-80; prepared in cooperation with Morse Laboratories, Inc.; CDL:236662-A)

MRID	CITATION
00020890	Uniroyal Chemical (1978) Vitavax®-25DB; Barley. (Unpublished study received Apr 25, 1979 under ND 79/8; prepared in cooperation with North Dakota State Univ. and Washington State Univ., Regional Cereal Disease Laboratory; CDL:238250-C)
00020888	Uniroyal Chemical (1978) Vitavax®-25DB; Oats. (Unpublished study received Apr 25, 1979 under ND 79/8; prepared in cooperation with North Dakota State Univ., South Dakota State Univ. and Univ. of Wisconsin, Dept. of Plant Pathology; CDL:238250-A)
00005867	Uniroyal Chemical (1979) Residue Chemistry: [Vitavax®]. (Unpublished study received Feb 6, 1979 under 400-EX-58; prepared in cooperation with Morse Laboratories, Inc.; CDL:237329-J)
00003233	Uniroyal Chemical (1979) Residues in PPM: Vitavax®. (Unpublished study received Mar 20, 1979 under 400-80; prepared in cooperation with Morse Laboratories, Inc.; CDL:098014-E)
00024917	Uniroyal Chemical (1979) Summary: [Vitavax®]. (Unpublished study received Oct 18, 1979 under 400-112; prepared in cooperation with Morse Laboratories, Inc.; CDL:241184-A)
00024922	Uniroyal Chemical (1979) Toxicology Vitavax® 30C. Summary of studies 240917-B and 240917-C. (Unpublished study received Aug 24, 1979 under 400-124; CDL:240917-A)
00005855	Uniroyal Chemical (1979) Toxicology: Vitavax® 30C. (Unpublished study received Feb 28, 1979 under 400-124; CDL:237477-A)
00032175	Uniroyal Chemical (1979) Vitavax® Fungicide: Uniroyal. (Technical data sheet; also In unpublished submission received Jun 3, 1980 under OK 80/9; submitted by Oklahoma, Dept. of Agriculture, Oklahoma City, Okla.; CDL:242612-A)
00030655	Uniroyal Chemical (1980) Residues in PPM. (Unpublished study received Jun 28, 1980 under NC 80/14; submitted by state of North Carolina, Dept. of Agriculture, Pesticide and Plant Protection Div. for Uniroyal; CDL:242701-A)
00005869	Uniroyal, Incorporated (1977) Technical Data Sheet: Uniroyal Vitavax® Fungicide. (Unpublished study received Apr 21, 1977 under 400-107; CDL:237180-A)
00021613	Uniroyal, Incorporated (1977) Vitavax®: Systemic Fungicide. (Unpublished study received Sep 18, 1967 under 400-EX-30; CDL:123433-K)
00021612	Uniroyal, Incorporated (1967) Bibliography of Vitavax® and Plantvax Systemic Fungicides. (Unpublished study received Jun 13, 1967 under 400-EX-30; CDL:123433-J)
00025463	Uniroyal, Incorporated (1970) Summary of Work on Plant Residues from Vitavax® Seed Treatments. (Unpublished study received Jan 12, 1970 under OF0939; CDL:091604-A)

MRID	CITATION
00025480	Uniroyal, Incorporated (1971) Composition of Vitavax®-Technical Grade. (Unpublished study received Jun 17, 1974 under 5F1525; CDL:094043-D)
00025483	Uniroyal, Incorporated (1974) Summary of Corn and Rice Residue Data. (Unpublished study received Jun 17, 1974 under 5F1525; prepared in cooperation with Morse Laboratories, Inc. and State Univ. College of New York--Oswego, Lake Ontario Environmental Laboratory; CDL:094043-H)
00003100	United States Rubber Company (1977) Vitavax® and Plantvax®: Systemic Fungicides: Agricultural Chemicals: Technical Data Sheet: Experimental Fungicide. (Unpublished study received Jun 13, 1967 under unknown admin. no.; CDL:107975-A)
05005921	Unverricht, A.; Schuette, H.R.; Banasiak, L.; Grahl, R.; Dedek, W. (1978) Synthesen radioaktiv markierter Verbindungen; 1) Darstellung von 5,6-Dihydro-2-methyl-1,4-oxathiin-3-carbonsaeureanilid-phenyl-U-14C (Carboxin), Carboxin-sulfoxid- und Carboxin-sulfon-phenyl-U-14C. [Syntheses of radioactively labeled compounds; 1) presentation of 5,6-dihydro-2-methyl-1,4-oxathiino-3-carboxylic acid-anilide-phenyl-U-14C (carboxin), carboxin-sulfoxide- and carboxin-sulfonphenyl-U-14C.] [letter]. Zeitschrift fuer Chemie 18(1):21-22.
05013071	Venkata Ram, C.S. (1977) Laboratory and field evaluation of carboxin and oxycarboxin against <u>Exobasidium vexans</u> on tea. Indian Phytopathology 30(4):449-455.
05003851	Verzar-Petri, G.; Haggag, M.Y. (1976) Gas liquid chromatographic method for detecting and measuring tropane alkaloids and Vitavax fungicide in the same extract, of <u>Datura innoxia</u> Mill. Herba Hungarica 15(1):87-96.
00003339	Von Schmeling, B.; Kulka, M. (1968) Systematic fungicide 2,3-Dihydro-5-carboxanilido-6-methyl-1,4-oxathiin. Pages 1-2, In Fungicidal Control of Smut Diseases of Cereals. Compiled by J.G. Moseman. Beltsville, Md.: U.S. Dept. of Agriculture. (U.S. Agricultural Research Service, Crops Research Div., CR 42-68; also In unpublished submission including glossary, received Mar 25, 1969 under 9G0819; submitted by Uniroyal Chemical, Bethany, Conn.; CDL:093520-AC)
05004276	Von Schmeling, B.; Kulka, M.; Thiara, D.S.; Harrison, W.A., inventors; United States Rubber Company, assignee (1966) Control of plant diseases. U.S. patent 3,249,499. May 3. 6 p. U.S. Cl. 167-33.
05004275	Von Schmeling, B.; Kulka, M., inventors; Uniroyal, assignee (1969) Stimulation of plant growth. U.S. patent 3,454,391. Jul 8. 3 p. Int. Cl. A Oln 5/00, 21/02; U.S. Cl. 71-77.

MRID

CITATION

- 00003059 Weir, R.J. (1966) Final Report: Acute Toxicity in Aqueous Exposure to Bluegill Sunfish: Project No. 798-100. (Unpublished study received Dec 14, 1966 under 400-EX-27; prepared by Hazleton Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:123429-B)
- 00003037 Weir, R.J. (1967) Final Report: Acute Dietary Toxicity--Bobwhite Quail: Project No. 798-100. (Unpublished study received Jun 14, 1969 under 9G0819; prepared by Hazleton Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091420-O)
- 00010348 Weir, R.J. (1967) Final Report: Acute Dietary Toxicity--Bobwhite Quail: Project No. 798-100. (Unpublished study received Nov 16, 1967 under 400-75; prepared by Hazleton Laboratories, Inc. for United States Rubber Co., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:132110-B)
- 00010347 Weir, R.J. (1967) Final Report: Acute Toxicity in Aqueous Exposure to Rainbow Trout: Project No. 798-100. (Unpublished study received Dec 11, 1967 under 400-75; prepared by Hazleton Laboratories, Inc. for United States Rubber Co., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:132110-A)
- 00021615 Weir, R.J. (1967) Final Report: Acute Toxicity in Aqueous Exposure to Rainbow Trout: Project No. 798-100. (Unpublished study received Jun 13, 1967 under 400-EX-30; prepared by Hazleton Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:123433-R)
- 00003036 Weir, R.J. (1967) Final Report: Acute Toxicity in Aqueous Exposure to Rainbow Trout: Project No. 798-100. (Unpublished study received Jun 14, 1969 under 9G0819; prepared by Hazleton Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:091420-M)
- 00021627 Weir, R.J. (1967) One-Year Progress Report: Two-Year Dietary Administration--Dogs: Project No. 798-103. (Unpublished study received Jul 19, 1968 under 400-80; prepared by Hazleton Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:003258-L)
- 00021628 Weir, R.J. (1967) 12-Month Progress Report: 24-Month Dietary Administration--Albino Rats: Project No. 798-102. (Unpublished study received Jul 19, 1968 under 400-80; prepared by Hazleton Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:003258-M)
- 00029175 Weir, R.J.; Hopkins, M. (1967) Final Report: Acute Dietary Toxicity--Bobwhite Quail: Project No. 798-100. (Unpublished study received Dec 12, 1967 under 400-49; prepared by Hazleton Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:009021-H)

MRID

CITATION

- 00029174 Weir, R.J.; Rathbun, F.; Hopkins, M. (1967) Acute Toxicity in Aqueous Exposure to Bluegill Sunfish and Rainbow Trout: Project No. 798-100. Final rept. (Unpublished study received Dec 12, 1967 under 400-49; prepared by Hazleton Laboratories, Inc., submitted by Uniroyal Chemical, Bethany, Conn.; CDL:009021-G)
- 05006394 White, R.P.; Johnston, H.W. (1972) Possible influences of Vitaflo on the growth of corn. Canadian Journal of Plant Science 52(6):1078-1080.
- 05003776 Yip, G. (1971) Report on herbicides, plant growth regulators, and fungicides. Journal of the Association of Official Analytical Chemists 54(2):327-330.
- 05003775 Yip, G. (1972) Report on fungicides, herbicides, and plant growth regulators. Journal of the Association of Official Analytical Chemists 55(2):287-289.
- 05008662 Zadoks, J.C.; Kodde, A.; Hoogkamer, W. (1969) The effect of derivatives of 1,4-oxathiin on Puccinia horiana in Chrysanthemum morifolium. Netherlands Journal of Plant Pathology 75(1/2):193-196.
- 05006396 Zimmer, D.E. (1967) Efficacy of 1,4-oxathiin fungicides for control of seedling safflower rust. Plant Disease Reporter 51(7):586-588.